



HAL
open science

The p76 Rb and p100 Rb truncated forms of the Rb protein exert antagonistic roles on cell death regulation in human cell lines

Nathalie Le Floch, Vincent Rincheval, Ioana Ferecatu, Rahamata Ali-Boina, Flore Renaud, Bernard Mignotte, Jean-Luc Vayssière

► To cite this version:

Nathalie Le Floch, Vincent Rincheval, Ioana Ferecatu, Rahamata Ali-Boina, Flore Renaud, et al.. The p76 Rb and p100 Rb truncated forms of the Rb protein exert antagonistic roles on cell death regulation in human cell lines. *Biochemical and Biophysical Research Communications*, 2010, 399 (2), pp.173-178. 10.1016/j.bbrc.2010.07.041 . hal-02977708

HAL Id: hal-02977708

<https://hal.uvsq.fr/hal-02977708>

Submitted on 26 Oct 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The p76^{Rb} and p100^{Rb} truncated forms of the Rb protein exert antagonistic roles on cell death regulation in human cell lines

Nathalie Le Floch ^{a,1}, Vincent Rincheval ^{a,1,*}, Ioana Ferecatu ^a, Rahamata Ali-Boina ^a, Flore Renaud ^a, Bernard Mignotte ^a and Jean-Luc Vayssière ^a

^a Université de Versailles / St Quentin-en-Yvelines, Laboratoire de génétique et biologie cellulaire / Ecole Pratique des Hautes Etudes, Laboratoire de génétique moléculaire et physiologique, 45 avenue des Etats-Unis, 78035 Versailles cedex, France.

* Corresponding author. Address: UVSQ, LGBC, 45 avenue des Etats-Unis, 78035 Versailles cedex, France.

Fax: (+33) 0139253655

E-mail address: vincent.rincheval@uvsq.fr

¹ Both authors contributed equally to this work

Abbreviations: eto, etoposide; E/TNF, emetine plus TNF- α ; Rb, retinoblastoma protein; TNF, tumor necrosis factor; CMX-ros, chloromethyl-X-rosamine; $\Delta\Psi_m$: mitochondrial membrane potential.

Footnotes:

Ioana Ferecatu present address is Institut de Chimie des Substances Naturelles, CNRS-UPR 2301, équipe 34, 1 avenue de la Terrasse, 91190 Gif-sur-Yvette, France.

Rahamata Ali-Boina present address is Laboratoire d'immunologie et immunothérapie des cancers de l'EPHE, Unité Inserm 866, Faculté de Médecine, 7 Bd Jeanne d'Arc, BP 87900 21079 Dijon cedex, France.

ABSTRACT

Several caspase-cleaved forms of the retinoblastoma protein have been described. Here, we compared the effect of full-length Rb versus the truncated p76^{Rb} and p100^{Rb} proteins on cell death regulation in five human cell lines. Interestingly, we observed that p76^{Rb} triggers cell death in all tested cell lines and that p100^{Rb} protects two cell lines against etoposide or TNF- α -induced cell death, whereas full-length Rb has no apoptotic effect. These results show that truncated forms of Rb can have specific activities in the regulation of cell death. They also suggest that caspase-cleavage of Rb should not be simply assimilated to a degradation process. Finally, we show that cell death induced by p76^{Rb} is Bax-dependent and is diminished by Bcl-2 overexpression or by caspase inhibition and that p100^{Rb} could inhibit cell death by decreasing both p53 stability and caspase activity.

Keywords: Retinoblastoma protein; Rb cleavage; caspase; apoptosis, transfection, GFP.

Introduction

The retinoblastoma protein (Rb) was the first identified tumour suppressor protein. Its loss of function is linked to the development of many human cancers. It has been shown that overexpression of Rb can induce growth arrest in a major G1 checkpoint blocking S-phase entry. The Rb protein governs cell cycle progression by controlling the expression of E2F-dependent genes in two ways: by directly binding and blocking the activation domain of E2F proteins or by active repression through chromatin remodelling.

Evidences for the importance of Rb in cellular differentiation and cell death come from studies of Rb knock-out mice, where the disruption of the Rb gene causes death by day 14 of gestation associated with defects in the development of the haematopoietic system, as well as in the central and peripheral nervous systems [1-3]. Interestingly, abolition of Rb function results in apoptosis involving both p53-dependent and -independent pathways [4, 5]. The transcription factor E2F-1 was proposed to be one of the possible links between Rb and p53. Indeed, free E2F-1 can transcriptionally activate p19^{ARF}, which stabilizes p53 by inhibiting the function of its regulator MDM2 [6]. Overexpression of Rb, which leads to a sequestration of E2F-1, can therefore inhibit apoptosis induced by p53.

A pro-apoptotic function of Rb has also been observed but is less well documented. Most of these studies show a direct role of Rb in the induction of apoptosis either by transcriptional activation or repression [7-9]. One model describes that Rb interacts with p53 and MDM2 in a trimeric complex that stabilizes both p53 and MDM2, enhancing p53-mediated transrepression which can trigger apoptosis [10]. More recently, two distinct Rb-E2F-1 complexes have been described, one of which activates transcription from pro-apoptotic genes to directly promote cell death [11]. These data suggest that the pro-apoptotic activities of Rb might be related to the complexes that it could form with its partners. Indeed,

our studies in drosophila have shown that the capacity of the drosophila homolog of Rb to be pro-apoptotic depends on the proliferation status of the cells in which it is expressed [12].

During apoptosis, Rb was shown to be cleaved by caspases, either at a C-terminal caspase-3-like consensus site generating a p100^{Rb} form, lacking the last 42 amino-acids [13-15], or at an internal cleavage site producing two fragments of 68 and 48 kDa [16].

Transgenic mice containing a germline mutation (*RB-MI*) at the caspase cleavage site generating p100^{Rb} were created. These mice exhibit resistance to apoptosis in selective tissues in response to specific death stimuli [17] and it was shown that this resistance involves a defect in Bid cleavage [18]. Recently, we identified the previously undescribed cleavage of Rb, at a LExD site, generating the p76^{Rb} form, which antagonizes p53-induced apoptosis in rat embryo fibroblasts [19].

In this article, we compared the effect of full-length Rb versus p76^{Rb} and p100^{Rb}, in several human cell lines. Interestingly, we report that although full length Rb does not have any apoptotic effect, p76^{Rb} triggers cell death in the absence of any other apoptosis inducers in all the tested cell lines and p100^{Rb} protects two cell lines against etoposide or TNF- α induced cell death. These results show that truncated forms of Rb can have specific activities in the regulation of cell death if compared to the full-length protein. Finally, we show on the one hand that p76^{Rb}-induced cell death is Bax-dependent and can be diminished by Bcl-2 overexpression or caspase inhibition and on the other hand that p100^{Rb} could inhibit cell death by decreasing p53 protein level and caspase activity.

Materials and methods

Reagents. Emetine (E), mouse tumour necrosis factor- α (TNF- α) and etoposide (eto) were from Sigma. Chloromethyl-X-rosamine (CMX-ros) was from Molecular Probes. z-Val-Ala-DL-Asp-Fluoromethylketone (zVAD) was from Bachem. Stock solutions were prepared as follows: emetine and TNF- α in serum free medium, etoposide and CMX-ros in dimethyl sulfoxide (DMSO), zVAD in methanol. All stock solutions were stored at -20°C .

Cell lines, cell culture and induction of cell death. We used the following human cell lines: HT1080, HEK293T, SAOS-2, TC7 and HT29. All cell lines were cultured at 37°C in a humidified atmosphere containing 5% CO_2 and in DMEM/F12 medium supplemented with 10% fetal bovine serum together with penicillin (100 $\mu\text{g}/\text{ml}$), streptomycin (100 U/ml) and glutamax (1% v/v) from Invitrogen. Cell death was induced by the addition to exponentially growing adherent cells of etoposide or TNF- α plus emetine during 24h. The concentrations of apoptosis inducers were determined to induce between 30% and 60% cell death after a 24 h incubation. The following doses were used: 125 $\mu\text{g}/\text{ml}$ etoposide for HEK293T, 30 $\mu\text{g}/\text{ml}$ etoposide or 0.4 $\mu\text{g}/\text{ml}$ emetine plus 2.5 ng/ml TNF- α for HT1080, 1.5 $\mu\text{g}/\text{ml}$ emetine plus 12.5 ng/ml TNF- α for SAOS-2, HT29 and TC7. zVAD was used at 100 μM .

Plasmids. The GFP-Rb (pEGFPC1-Rb) construct was previously described [20] with Rb cDNA of human origin. cDNAs corresponding to truncated p100^{Rb} and p76^{Rb} were subcloned into pGEM-T vectors (Promega) after PCR amplification. PCR was performed on human wild type Rb coding sequence (NM_000321) contained in pEGFPC1-Rb plasmid.

Primers used for PCR amplification are 5'-

TCTCGAGCTATGCCGCCCAAACCCCCCGAAAA-3' and 5'-

GAAGCTTTCAATCTGCTTCATCTGATCCTTCA-3' for p100Rb; 5'-

TCTCGAGCTATGACAAGAATTATTGAAGTTC-3' and 5'-

GAAGCTTTCATTTCTCTTCCTTGTTTGAGGT-3' for p76^{Rb} (italic: XhoI or HindIII

restriction sites, bold: start or stop codon introduced when necessary, underlined: specific Rb

sequence). PCR was performed using standard methods with DyNAzyme™ EXT DNA

polymerase (Finnzymes), performing 5 PCR cycles at 37°C (denaturation at 94°C for 30s,

annealing at 37°C for 30 s and extension at 72°C for 3 min) followed by 25 PCR cycles at

55°C (denaturation at 94°C for 30 s, annealing at 55°C for 30 s and extension at 72°C for 3

min), except for p100^{Rb} amplification which required 30 cycles at 50°C instead of 25 cycles at

55°C. PCR products were extracted from agarose gel with JETsorb kit and ligated into

pGEM-T vector using T4-DNA-ligase according to manufacturer's instructions (pGEM-T kit,

Promega). pGEM-T-p100^{Rb} and -p76^{Rb} vectors were next digested using XhoI and HindIII

enzymes and purified p100^{Rb} and p76^{Rb} fragments were ligated into XhoI/HindIII linearized

pEGFPC1 vector (Invitrogen). Fusion cDNA sequences were checked by sequencing both

strands using appropriate primers (MWG biotech). The use of pEGFPC1 vector conferred a

GFP tag at the N-terminus of Rb, p76^{Rb} and p100^{Rb}.

Transfection. The nucleofection technique was performed according to the manufacturers' instructions (Amaxa) and cells were transfected with the following combinations of nucleofection solutions and programs: HEK293T (L / A-020), SAOS-2 (V / D-024), TC7 (T / T-030), HT29 (R / W-017). The HT1080 cells were transfected with the lipofectamin 2000 technique according to the manufacturer's instructions (Invitrogen). Induction of cell death was performed 24 h after the beginning of transfection.

SiRNA treatments. Cells were seeded 24 h before transfection with control non silencing (QIAGEN) or Bax siRNA (Hs BAX 10 HP validated siRNA, QIAGEN). Lipofection was made using HiPerfect reagent (QIAGEN) and 25 nM siRNA according to the manufacturer's instructions. The cells were incubated with siRNA for 24h at 37°C before transfection with GFP-Rb or GFP-p76^{Rb} encoding vectors, as described above. Experiments (protein extraction for western-blot analysis and flow cytometry analysis) were carried out 24h to 48h later.

Flow cytometry. Flow cytometric measurements were performed using a XL3C flow cytometer (Beckman-Coulter). Cells expressing GFP-tagged proteins were identified by the emission of green fluorescence (525 nm) and mitochondrial membrane potential ($\Delta\Psi_m$) was assessed by the retention of CMX-ros (620 nm). This fluorochrome, at low doses, specifically accumulates into the mitochondria depending on $\Delta\Psi_m$. As a drop in $\Delta\Psi_m$ characterises all types of cell death [21], cell viability was determined by measuring the percentage of cells with low CMX-ros fluorescence. Cell death was also checked by assessing the decrease in cell size / forward scatter that reflects cell shrinkage during cell death. These measurements were performed in GFP positive cells in order to characterize the effect of exogenous proteins on cell viability. The expression level of the fusion proteins was systematically assessed by measuring GFP fluorescence intensity. Briefly, the cells were trypsinized and pooled with their corresponding medium and centrifuged (200g, 5 min). Cells were then resuspended in complete medium with 150 μ M CMX-ros and analyzed under FACS after 30 min incubation at 37°C.

Western blot analysis. Western blotting was performed as previously described [22]. The primary antibodies used were mouse monoclonal anti-p53 (Pab 122, gift from Dr. E.

May, IRSC, Villejuif, France), anti-MDM2 (SMP40, Santa Cruz), anti-enolase (gift from N. Lamande, College de France, Paris), anti-cleaved PARP (Asp214, BD Pharmingen), anti-Rb (G3-245, BD Pharmingen), rabbit polyclonal anti-p53-P (Ser-15, Santa Cruz), anti-caspase 9 (Cell Signaling) and anti-cleaved caspase-3 (Asp175, Cell Signaling) as well as goat polyclonal anti-PUMA- α (N-19, 1:200, Santa Cruz). Secondary antibodies (peroxidase-conjugated) were anti-mouse, anti-rabbit or anti-goat immunoglobulin (Biosystem). Immunoreactive spots were detected by ECL or ECLplus kits depending on signal intensity (Amersham).

Results

Our aim was to characterise potential roles of p100^{Rb} and p76^{Rb} in several human cell lines. To perform this study, we have used GFP-tagged proteins in order to assess cell viability only in cultured cells that were successfully transfected. We have also used cell lines with distinct p53 status (summarized in Fig. 1A) in order to test if p100^{Rb} and p76^{Rb} activities depended on p53. Each cell line was transfected with vectors encoding the three forms of Rb (full length Rb, p100^{Rb} and p76^{Rb}) and 24 hours later, the cells were incubated or not with apoptosis inducers (emetine + TNF- α (E/TNF) or etoposide) during another 24 hours. The cells were then subjected to flow cytometric analysis to evaluate their viability (see Materials and Methods). Etoposide is a topoisomerase II inhibitor which can trigger the intrinsic pathway of apoptosis [23] whereas TNF- α is a cytokine which can trigger the extrinsic pathway of cell death [24]. TNF- α was used in combination with emetine, an inhibitor of protein synthesis which (at non toxic concentrations) potentiates the pro-apoptotic effect of this cytokine [25, 26]. HT29, TC7 and SAOS-2 were not sensitive to etoposide (probably because of their deficient p53 status) whereas HEK293T cells were insensitive to E/TNF (data not shown). The HT1080 cell line was the only one to be sensitive to inducers of both intrinsic and extrinsic apoptotic pathways. After drugs incubation, cellular viability was measured on GFP positive cells by characterizing the mitochondrial membrane potential ($\Delta\Psi_m$) which has been shown to decrease during apoptosis, autophagy or necrosis [21]. Cell shrinkage was also assessed by flow cytometry to confirm the results (data not shown).

Fig. 1A shows that the full-length Rb protein has no significant effect (if compared to GFP alone) on cellular viability, in the presence or in the absence of apoptosis inducers, in all tested cell lines. This is also the case for p100^{Rb} in the absence of apoptosis inducers. Interestingly, we show here that p76^{Rb} significantly induces cell death on its own in the five

tested cell lines (compare GFP or GFP-Rb to GFP-p76^{Rb}). This suggests that the effect of p76^{Rb} is independent of the status of p53. p76^{Rb} also increases the level of cell death in the presence of apoptosis inducers, except in HT29 cells. We have also observed that p100^{Rb} inhibits apoptosis in HT1080 + etoposide and SAOS-2 + E/TNF (compare GFP or GFP-Rb to GFP-p100^{Rb}). The results obtained with the three forms of Rb are very interesting since they show that the truncated forms of Rb could have new and specific activities. Fig. 2B shows that the effects of these proteins are not related to a difference in their expression levels, indicated by the measurements of GFP fluorescence intensities. The fact that p76^{Rb} appears to be a little less expressed may be due to its specific toxicity.

In order to better understand how p76^{Rb} and p100^{Rb} regulate cell death, we tried to characterize several aspects of their mechanism of action. Fig. 2A shows that the broad spectrum caspase inhibitor zVAD significantly inhibits p76^{Rb} induced cell death in TC7 and HT29 cell lines, demonstrating that this cell death can be caspase-dependent. We have also used a variant of the SAOS-2 cell line conditionally overexpressing *bcl-2* (Tet off system from Clontech) and showed that Bcl-2 significantly inhibits p76^{Rb} induced cell death (Fig. 2B). Finally, we have tested the effect of Bax silencing on p76^{Rb} induced cell death in HT1080 cells. Fig. 2C (left panel) shows the efficiency of the siRNA treatment and points out that p76^{Rb} is able to trigger the accumulation of the pro-apoptotic Bax protein. Interestingly, it can also be observed from the corresponding analysis of cellular viability (right panel) that Bax siRNA decreases p76^{Rb} induced cell death, showing that it is Bax-dependent. Taken together, these results strongly suggest that GFP-p76^{Rb} triggers cell death by inducing the mitochondrial (intrinsic) pathway of apoptosis.

Next, we have investigated the mechanism underlying the anti-apoptotic role of p100^{Rb} in the HT1080 cells incubated with etoposide. We show in Fig. 3 that p100^{Rb} inhibits the accumulation of both total and the active p53 (phosphorylated on serine 15), triggered by

etoposide. Interestingly, this phenomenon occurs very early (between 5 and 8h of incubation) and is no more visible after 16h (coinciding with a decrease in p100^{Rb} amount). Nevertheless, it seems to have consequences on the p53 transcriptional target and pro-apoptotic protein PUMA since its accumulation is specifically inhibited afterward. Interestingly, p100^{Rb} also seems to inhibit processing of caspase-9 and caspase-3 as well as caspase-3 activity, visualised by PARP cleavage, after 16 h of incubation. It is thus tempting to hypothesize that the upstream events seen with p53 stability are responsible for the caspase inhibition and the anti-apoptotic activity of p100^{Rb}.

Discussion

Here, we have shown that p76^{Rb} and p100^{Rb} have specific roles on cell death regulation when compared to full-length Rb. In particular, we observed that p76^{Rb} induces cell death in the five tested human cell lines, probably via the activation of a mitochondrial-dependent apoptotic pathway. We have also observed an anti-apoptotic activity of p100^{Rb} in two cell lines. We have shown in the HT1080 cell line that p100^{Rb} interferes with the stability and activity of the p53 protein whereas this mechanism appears impossible in SAOS-2 cells, since they are p53-deficient. This matter of fact strongly suggests that p100^{Rb} could act at multiple levels to inhibit cell death. Indeed, if we consider that p100^{Rb} inhibits both p53 and caspase activity, we can hypothesize that the inhibition of caspase activity could be partly independent from the effect on p53. Concerning the interplay between p100^{Rb} and p53, two partners of Rb were already described to be involved in the regulation of p53 stability: MDM2 and E2F-1 [6, 27]. First, it has been reported that p100^{Rb} exhibit an enhanced E2F binding [14], suggesting that p100^{Rb} could inhibit E2F-1-dependent expression of p19^{ARF} and consequently p53 stability [6]. Second, it has been shown that, in contrast to full-length Rb, p100^{Rb} cannot interact with MDM2 [13]. Thus, the C-terminal cleavage of Rb could prevent the formation of the Rb-MDM2-p53 trimeric complex which is known to stabilize p53 [10]. In conclusion, implications of both E2F-1 and MDM2 proteins could be in agreement with the anti-apoptotic role of p100^{Rb}.

Finding that p76^{Rb} significantly induces cell death in all tested human cell lines is interesting since it suggests that this effect is not cell type dependent. The effect of p76^{Rb} is probably independent from p53 since it occurs in both wild type (HT1080) and p53-deficient (TC7, HT29, SAOS-2) cell lines (see Fig. 1A). However, we cannot exclude that several pathways could be triggered by p76^{Rb}, some pathways being p53-dependent or independent. Finally, by using GFP-tagged proteins, we have observed that full-length Rb, p100^{Rb} and

p76^{Rb} are located into the nucleus (data not shown). The fact that these three forms of Rb exhibit distinct activities suggests that they interact with different partners into this compartment to regulate the mitochondrial-dependent pathway of apoptosis. Thus a differential regulation of gene expression is probably central to the activity of the p76^{Rb} and p100^{Rb} truncated forms of the retinoblastoma protein. However the specific partners and the precise mechanism of this regulation remain to be determined.

The role of full length Rb in cell death regulation is complex since Rb was shown to exert opposite effects, depending on the cellular context. Our study suggests that such a complexity could exist for truncated forms of Rb since we have shown a few years ago that p76^{Rb} can inhibit a p53-dependent cell death in immortalized or primary rodent fibroblasts [19]. The pro-apoptotic effect of p76^{Rb} was first found in the human HEK293T cell line in which p53 may be inactivated by the presence of two viral oncoproteins: E1B and LT. Thus we have hypothesized that a difference in p53 activity could be responsible for the discrepancies between our two studies. However, we have tested the effect of p76^{Rb} on human cell lines with distinct p53 status, and found that p76^{Rb} was intrinsically and systematically pro-apoptotic. Since this new activity of p76^{Rb} was a general observation in human cell lines, we now postulate that cell death regulations by the Rb pathway might be different between rodent and human cells, explaining the opposite roles of p76^{Rb} in our two independent studies. Such an hypothesis would be in agreement with several of our previous studies, which showed that caspase inhibition can paradoxically accelerate cell death in rodent cells [19, 28-30] whereas we have never retrieved this effect in human cells. Finally, we are currently using *Drosophila melanogaster* in our laboratory as an *in vivo* model to study the role of truncated forms of the Rb homolog. Our results suggest that p76^{Rb} accelerates cell death in specific tissues if compared to full length Rb (unpublished data), reinforcing the significance of the present study.

Acknowledgments

This work was supported in part by grants from the Association pour la Recherche contre le Cancer (#3819) and from the Ligue Nationale Contre le Cancer. We thank Simon Heath for his critical reading of the manuscript. We are grateful to Christopher J. Hutchison who has provided the pEGFPC1-Rb plasmid.

References

- [1] E. Y. Lee, C. Y. Chang, N. Hu, Y. C. Wang, C. C. Lai, K. Herrup, W. H. Lee, and A. Bradley, Mice deficient for Rb are nonviable and show defects in neurogenesis and haematopoiesis, *Nature* 359 (1992) 288-294.
- [2] T. Jacks, A. Fazeli, E. M. Schmitt, R. T. Bronson, M. A. Goodell, and R. A. Weinberg, Effects of an Rb mutation in the mouse, *Nature* 359 (1992) 295-300.
- [3] A. R. Clarke, E. R. Maandag, M. van Roon, N. M. van der Lugt, M. van der Valk, M. L. Hooper, A. Berns, and H. te Riele, Requirement for a functional Rb-1 gene in murine development, *Nature* 359 (1992) 328-330.
- [4] S. D. Morgenbesser, B. O. Williams, T. Jacks, and R. A. DePinho, p53-dependent apoptosis produced by Rb-deficiency in the developing mouse lens, *Nature* 371 (1994) 72-74.
- [5] K. F. Macleod, Y. Hu, and T. Jacks, Loss of Rb activates both p53-dependent and independent cell death pathways in the developing mouse nervous system, *Embo J* 15 (1996) 6178-6188.
- [6] J. Pomerantz, N. Schreiber-Agus, N. J. Liegeois, A. Silverman, L. Alland, L. Chin, J. Potes, K. Chen, I. Orlow, H. W. Lee, C. Cordon-Cardo, and R. A. DePinho, The Ink4a tumor suppressor gene product, p19Arf, interacts with MDM2 and neutralizes MDM2's inhibition of p53, *Cell* 92 (1998) 713-723.
- [7] C. Bowen, S. Spiegel, and E. P. Gelmann, Radiation-induced apoptosis mediated by retinoblastoma protein, *Cancer Res* 58 (1998) 3275-3281.
- [8] X. Zhao, and M. L. Day, RB activation and repression of C-MYC transcription precede apoptosis of human prostate epithelial cells, *Urology* 57 (2001) 860-865.

- [9] C. Bowen, M. Birrer, and E. P. Gelmann, Retinoblastoma protein-mediated apoptosis after gamma-irradiation, *J Biol Chem* 277 (2002) 44969-44979.
- [10] J. K. Hsieh, F. S. Chan, D. J. O'Connor, S. Mittnacht, S. Zhong, and X. Lu, RB regulates the stability and the apoptotic function of p53 via MDM2, *Mol Cell* 3 (1999) 181-193.
- [11] A. Ianari, T. Natale, E. Calo, E. Ferretti, E. Alesse, I. Screpanti, K. Haigis, A. Gulino, and J. A. Lees, Proapoptotic function of the retinoblastoma tumor suppressor protein, *Cancer Cell* 15 (2009) 184-194.
- [12] C. Milet, A. Rincheval-Arnold, B. Mignotte, and I. Guenal, The *Drosophila* retinoblastoma protein induces apoptosis in proliferating but not in post-mitotic cells, *Cell Cycle* 9 (2010) 97-103.
- [13] R. U. Janicke, P. A. Walker, X. Y. Lin, and A. G. Porter, Specific cleavage of the retinoblastoma protein by an ICE-like protease in apoptosis, *Embo J* 15 (1996) 6969-6978.
- [14] W. D. Chen, G. A. Otterson, S. Lipkowitz, S. N. Khleif, A. B. Coxon, and F. J. Kaye, Apoptosis is associated with cleavage of a 5 kDa fragment from RB which mimics dephosphorylation and modulates E2F binding, *Oncogene* 14 (1997) 1243-1248.
- [15] A. L. Boutillier, E. Trinh, and J. P. Loeffler, Caspase-dependent cleavage of the retinoblastoma protein is an early step in neuronal apoptosis, *Oncogene* 19 (2000) 2171-2178.
- [16] B. An, and Q. P. Dou, Cleavage of retinoblastoma protein during apoptosis: an interleukin 1 beta-converting enzyme-like protease as candidate, *Cancer Res* 56 (1996) 438-442.

- [17] B. N. Chau, H. L. Borges, T. T. Chen, A. Masselli, I. C. Hunton, and J. Y. Wang, Signal-dependent protection from apoptosis in mice expressing caspase-resistant Rb, *Nat Cell Biol* 4 (2002) 757-765.
- [18] X. Huang, A. Masselli, S. M. Frisch, I. C. Hunton, Y. Jiang, and J. Y. Wang, Blockade of tumor necrosis factor-induced Bid cleavage by caspase-resistant Rb, *J Biol Chem* 282 (2007) 29401-29413.
- [19] C. Lemaire, N. Godefroy, I. Costina-Parvu, V. Rincheval, F. Renaud, P. Trotot, S. Bouleau, B. Mignotte, and J. L. Vayssiere, Caspase-9 can antagonize p53-induced apoptosis by generating a p76(Rb) truncated form of Rb, *Oncogene* 24 (2005) 3297-3308.
- [20] E. Markiewicz, T. Dechat, R. Foisner, R. A. Quinlan, and C. J. Hutchison, Lamin A/C binding protein LAP2alpha is required for nuclear anchorage of retinoblastoma protein, *Mol Biol Cell* 13 (2002) 4401-4413.
- [21] J. J. Lemasters, A. L. Nieminen, T. Qian, L. C. Trost, S. P. Elmore, Y. Nishimura, R. A. Crowe, W. E. Cascio, C. A. Bradham, D. A. Brenner, and B. Herman, The mitochondrial permeability transition in cell death: a common mechanism in necrosis, apoptosis and autophagy, *Biochim Biophys Acta* 1366 (1998) 177-196.
- [22] S. Bouleau, I. Parvu-Ferecatu, A. Rodriguez-Enfedaque, V. Rincheval, H. Grimal, B. Mignotte, J. L. Vayssiere, and F. Renaud, Fibroblast Growth Factor 1 inhibits p53-dependent apoptosis in PC12 cells, *Apoptosis* 12 (2007) 1377-1387.
- [23] N. O. Karpinich, M. Tafani, R. J. Rothman, M. A. Russo, and J. L. Farber, The course of etoposide-induced apoptosis from damage to DNA and p53 activation to mitochondrial release of cytochrome c, *J Biol Chem* 277 (2002) 16547-16552.
- [24] A. Thorburn, Death receptor-induced cell killing, *Cell Signal* 16 (2004) 139-144.

- [25] C. Sidoti-de Fraisse, V. Rincheval, Y. Risler, B. Mignotte, and J. L. Vayssiere, TNF-alpha activates at least two apoptotic signaling cascades, *Oncogene* 17 (1998) 1639-1651.
- [26] A. Dumay, V. Rincheval, P. Trotot, B. Mignotte, and J. L. Vayssiere, The superoxide dismutase inhibitor diethyldithiocarbamate has antagonistic effects on apoptosis by triggering both cytochrome c release and caspase inhibition, *Free Radic Biol Med* 40 (2006) 1377-1390.
- [27] N. Godefroy, C. Lemaire, B. Mignotte, and J. L. Vayssiere, p53 and Retinoblastoma protein (pRb): a complex network of interactions, *Apoptosis* 11 (2006) 659-661.
- [28] V. Rincheval, F. Renaud, C. Lemaire, B. Mignotte, and J. L. Vayssiere, Inhibition of Bcl-2-dependent cell survival by a caspase inhibitor: a possible new pathway for Bcl-2 to regulate cell death, *FEBS Lett* 460 (1999) 203-206.
- [29] N. Godefroy, C. Lemaire, F. Renaud, V. Rincheval, S. Perez, I. Parvu-Ferecatu, B. Mignotte, and J. L. Vayssiere, p53 can promote mitochondria- and caspase-independent apoptosis, *Cell Death Differ* 11 (2004) 785-787.
- [30] N. Godefroy, S. Bouleau, G. Gruel, F. Renaud, V. Rincheval, B. Mignotte, D. Tronik-Le Roux, and J. L. Vayssiere, Transcriptional repression by p53 promotes a Bcl-2-insensitive and mitochondria-independent pathway of apoptosis, *Nucleic Acids Res* 32 (2004) 4480-4490.

Legends to figures

Fig. 1. Effect of GFP-p76^{Rb} and GFP-p100^{Rb} versus GFP-Rb and GFP on cellular viability in five human cell lines. (A) The four vectors encoding GFP, GFP-Rb, GFP-p76^{Rb} and GFP-p100^{Rb} were transfected in human cell lines with different p53 status (in HEK293T cells, two oncoproteins are present: E1B from the adenovirus and LT from SV40 which inhibit at least partially p53). The cells were incubated or not with apoptosis inducers (24h) and the cellular viability was determined by flow cytometric analysis (see Materials and Methods).

*Statistically significant difference in the viability of cells transfected with vectors were indicated (GFP-Rb versus GFP; GFP-p76^{Rb} or GFP-p100^{Rb} versus GFP-Rb). Statistical analysis was determined using Student's t test. $P < 0.05$ was accepted as significant. (B) The relative GFP intensity was measured in each experimental condition to assess the expression level of the fusion proteins.

Fig. 2. Regulation of p76^{Rb}-induced cell death. All cell lines used here were transfected with the GFP and GFP-p76 encoding vectors. The TC7 and HT29 cells were incubated with the caspase inhibitor zVAD (100 μ M) (A). The conditionally overexpressing *bcl-2* / SAOS-2 cell line (Tet off system, Clontech) was used to evaluate the effect of Bcl-2 (B). The HT1080 cells were treated or not with Bax siRNA and western blots of Bax and enolase were performed (C). In all cases, percentage of cell death was measured by flow cytometric analysis.

*Statistically significant difference in the viability of cells transfected with vectors were indicated (-zVAD versus +zVAD, -Bcl-2 versus +Bcl-2). Statistical analysis was determined using Student's t test. $P < 0.05$ was accepted as significant.

Fig. 3. Effect of p100^{Rb} on p53 stability and caspase activation. The HT1080 cells were transfected with the GFP and GFP-p100^{Rb} encoding vectors and were incubated 24 hours later for different times with etoposide. Western blots were performed to assess the role of p100^{Rb} on p53 accumulation and activity, as well as caspase activation and activity.

Figure 1






Figure 1

Figure 2

A



B



C




Figure 2

Figure 3




Figure 3