



## Reducing plastic production: Economic loss or environmental gain?

Mateo Cordier, Takuro Uehara, Bethany Jorgensen, Juan Baztan

### ► To cite this version:

Mateo Cordier, Takuro Uehara, Bethany Jorgensen, Juan Baztan. Reducing plastic production: Economic loss or environmental gain?. Cambridge Prisms: Plastics, 2024, pp.1-32. 10.1017/plc.2024.3 . hal-04430631

**HAL Id: hal-04430631**

**<https://hal.uvsq.fr/hal-04430631>**

Submitted on 1 Feb 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Reducing plastic production: Economic loss or environmental gain?

Mateo Cordier<sup>a,\*</sup>, Takuro Uehara<sup>b</sup>, Bethany Jorgensen<sup>c</sup> and Juan Baztan<sup>a</sup>

<sup>a</sup> Research Centre Cultures–Environnements–Arctique–Représentations–Climat (CEARC), Université de Versailles-Saint-Quentin-en-Yvelines, Université Paris-Saclay, 11 Boulevard d’Alember, 78280 Guyancourt, France. [mateo.cordier@uvsq.fr](mailto:mateo.cordier@uvsq.fr), [juan.baztan@uvsq.fr](mailto:juan.baztan@uvsq.fr)

<sup>b</sup> College of Policy Science, Ritsumeikan University, 2-150 Iwakura-Cho, Ibaraki City, 567-8570 Osaka, Japan. [takuro@fc.ritsumei.ac.jp](mailto:takuro@fc.ritsumei.ac.jp)

<sup>c</sup> Civic Ecology Lab, Department of Natural Resources, Cornell University, 144 East Ave., Ithaca, NY 14853 Ithaca, USA. [bj265@cornell.edu](mailto:bj265@cornell.edu).

\* Corresponding author: Mateo Cordier, CEARC-OVSQ, 11 boulevard d’Alember, 78280 Guyancourt, France. Email: [mateo.cordier@uvsq.fr](mailto:mateo.cordier@uvsq.fr)

## Abstract

We reviewed economic and environmental studies on global plastic pollution and we estimate the global cost of actions towards zero plastic pollution in all countries by 2040 to be US\$ 18.3-158.4 trillion (cost of a 47% reduction of plastic production included). If no actions are undertaken, we estimate the cost of damages caused by plastic pollution from 2016-2040 to be US\$ 13.7-281.8 trillion. These ranges suggest it is possible that the costs of inaction are significantly higher than those of action. Plastic product sales will also generate a global benefit in the form of incomes (salaries, dividends, etc.) estimated to be US\$ 38.0 trillion over 2016-2040 in the case of inaction, and US\$ 32.7-33.1 trillion in case of action. Calculating benefit minus costs provides the net benefits: US\$ –120.4-19.7 trillion in case of action and US\$ –243.8-24.3 trillion in case of inaction. Net benefit ranges suggest action and inaction will both be beneficial when considering the high estimates. However, the low estimates show net benefits might be negative, which suggests inaction might generate a net cost for society that will be twice the cost of action. Our estimates are preliminary (several cost and benefit data are lacking).

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI.  
10.1017/plc.2024.3

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

32

33 **Key words:** plastic debris, economic analysis, waste management, marine plastic pollution, coastal and  
34 ocean.

35

36

37

## 38 **Impact statement**

39 Lau et al. (2020) show that reducing plastic production and replacing plastics with alternative materials  
40 could reduce the production of plastics by 47% in 2040. This would reduce plastic pollution in terrestrial  
41 and aquatic ecosystems. Other interventions are also needed, such as cleanups in oceans, rivers,  
42 beaches and all terrestrial ecosystems. Interventions such as reusing old plastic products, improved  
43 collection, sorting, recycling and disposal of municipal solid plastic waste are also required in many  
44 countries. Implementing all these interventions globally, in theory, would allow the environmental  
45 target of zero plastic debris in the global ecosystem by 2040 to be met. This would cost between  
46 US\$ 18000 billion and US\$ 158000 billion, meaning the cost of action is between the GDP of China and  
47 1.6 times the world GDP. On the other hand, if we do nothing to address plastic pollution, the cost of  
48 global environmental damages (estimated to be US\$ 14000-282000 billion) could be significantly higher  
49 than the cost of taking actions to end plastic pollution. These actions, will certainly produce  
50 environmental gain. They might also produce an economic gain but this requires further research to  
51 reduce uncertainty margins and confirm inaction is substantially more expensive than action.

52

53

54

## 55 **Introduction**

56 Plastics represent a group of polymers including natural, semi-synthetic, or synthetic materials that are  
57 malleable and can be modeled into solid objects (Chen and Yan, 2020). Natural plastics such as horn,  
58 tortoiseshell, amber, rubber and shellac have been worked with since antiquity. However, the first  
59 synthetic plastic, Bakelite, is more recent and was invented by a Belgian chemist Leo Baekland in 1907  
60 (Science museum, 2019; Baekland, 1909). With the salient plastic virtues of low-cost, being lightweight,  
61 durable, odorless, and versatile, among others, a large and rapid expansion of plastic manufacturing  
62 started in the 1950s (Chen and Yan, 2020). In 1950, the annual production of plastic goods amounted to  
63 2 million metric tons (MMT) globally and by 2018, it surpassed 450 MMT (Law and Narayan, 2022;  
64 Geyer et al., 2017). This global market growth is projected to be driven in the future largely by increasing  
65 plastic use in the construction, automotive, and electrical and electronics industries (Grand View  
66 Research, 2022).

67 Scientists realized in the 2010s that a significant share of the massive amounts of plastics manufactured  
68 since 1950 had not been appropriately managed at the products' end of life (Geyer et al., 2017). Plastic

waste mismanagement explains why plastics are now found in the form of plastic debris in absolutely all ecosystems: on land and in the ocean, even in its deepest parts at 11 km depth in the Mariana trench (Chiba et al., 2018), and on all continents, even in Antarctica (Lacerda et al., 2019). Among all manufactured products, plastics are among the toughest to decay. The decomposition period of plastic waste in the environment is poorly understood but recent studies suggest it might range from decades to centuries and even several thousand years for several types of plastic products (Law and Narayan, 2022). The half-life of plastic products ranges, for example, from 4.2 years to more than 2500 years for plastic bags and from 12 years to more than 2500 years for plastic bottles. The half-life is defined as the time in which the plastic material loses 50% of its original mass through natural biodegradation in the environment, which depends on environmental conditions (Chamas et al., 2020). These estimations must, however, be considered cautiously as underlined in Ward and Reddy (2020). They show the extreme difficulty of estimating degradation times and defining what “plastic degradation” means.

Annual discards of inadequately managed plastic waste have been estimated by Lebreton and Andrady (2019), Lau et al. (2020), Cordier et al. (2021), and Yan et al. (2022). Annual discards have been increasing at the global scale, for example, from 23-91 MMT per year in 2010 to 36-115 MMT per year in 2020, and will probably multiply by 2-4 over the period of 2020-2060 (Figure S1 in Supplemental materials). Inadequately managed plastic waste is highly likely to be encountered in ecosystems since it includes littered plastic waste (directly thrown on the ground by individuals) and plastics for which waste treatment consists of collective discarding in waterways and marine areas or landfilling in open dumps, making it likely to enter terrestrial or marine ecosystems via inland waterways, wastewater outflows, storm drains, transport by wind or tides or leakages from open dumps and open uncontrolled landfills (Jambeck et al., 2015; Cordier et al., 2021; Lebreton and Andrady, 2019). These annual flows of inadequately managed plastic waste accumulate over time in the environment. Summing annual flows year after year gives the total amount of plastic accumulated since 1950, which passed from 444-2451 MMT in 2010 to 735-3373 MMT in 2020 and is forecast to be multiplied by 3-9 between 2020-2060 if no serious plastic pollution reduction strategies are undertaken in the coming years (Figure 1).

A portion of the globally accumulated discards of inadequately managed plastic waste since 1950 (Figure 1) leaks into the environment and accumulates in terrestrial and aquatic ecosystems (Figure 2). The massive amounts of plastic debris accumulated in ecosystems explain why marine scientists have detected plastic particles in a wide variety of marine organisms including mussels, oysters, shrimps, daphnia, turtles, sea birds, fish, etc. (Peng et al., 2020). Across all studies accounting for microplastics, the incidence rate of plastic ingested by fish was 26%. Over the last decade this incidence has doubled, increasing by 2.4% per year (Savoca et al., 2021). This presents serious threats to the health of marine animals, causing symptoms such as malnutrition, inflammation, chemical poisoning, growth thwarting, decrease of fecundity, and death due to damages at individual, organ, tissue, cell, and molecular levels (Peng et al., 2020). This means human health is also affected through seafood consumption. Plastic particles have been detected in human blood (Leslie et al., 2022) and in human placenta (Ragusa et al., 2021). Human health could be adversely affected stemming from both the exposure to chemicals contained in plastic components and from toxins that adsorb onto plastic debris from the surrounding seawater (Choy et al., 2019).

The accelerated accumulation of plastic debris in the environment since the 2000's raises three questions that can no longer be avoided: (i) should we clean terrestrial and aquatic ecosystems polluted with plastics; (ii) should we stop producing and consuming plastics to avoid future pollution; and (iii) is the cost of both options affordable and lower than the cost of inaction? The following sections help answer

these questions. Section 1 provides global estimations of the total amount of plastic debris accumulated in aquatic and terrestrial ecosystems. Section 2 presents strategies to reduce plastic contamination of ecosystems and the cost of action. Section 3 shows the global cost of the impacts that will result from plastic pollution in case of inaction from now to 2040. Section 4 provides a calculation of the net benefits (that is, benefits minus costs) earned from plastic sales. Section 5 discusses the results, compares the cost and net benefits of action and inaction, and concludes.

## 1. Global estimations of plastic debris accumulated in the ecosystems

The total amount of plastic accumulated in global terrestrial ecosystems since 1950 is estimated to be 320-629 MMT in 2020 and is forecast to multiply by 2.6 by 2040 (Figure 2, upper graph). In aquatic ecosystems, the global amount accumulated since 1950 is estimated to be 83-605 MMT in 2020 and is forecast to multiply by 1.5 or 2 by 2040 (Figure 2, lower graph).

To calculate some of the costs of plastic pollution reduction strategies (Section 2), it is important to distinguish the compartments of aquatic ecosystems where plastic debris accumulate since they require distinct removal and cleanup technologies. Global plastic accumulation in the oceans since 1950 is estimated to be 18-385 MMT in 2020 (Figure 3). Once it reaches the ocean, plastic debris may move to different parts of the marine environment. Data from the OECD (2022, p. 126) suggest that 87.8% of plastics reaching the global ocean are floating close to the ocean shoreline, 9.8% sink to the seabed, and 2.4% are transported offshore by marine currents and continue floating on the ocean surface (Figure 3). In rivers, the accumulation of floating plastics is estimated to be 18-45 MMT in 2020. For plastic debris sinking to riverbeds and lakebeds, accumulated amounts are estimated to be 46-114 MMT in 2020 (Figure 3).

## 2. Global cost of actions towards zero plastic debris in ecosystems by 2040

Plastic pollution reduction strategies can be organized into three categories (Lau et al., 2020; Cordier et al., 2019): (i) upstream preventive strategies designed to avoid plastics being produced (implemented at pre-consumption stages, e.g., reducing production and demand of plastics); (ii) mid-stream preventive strategies aimed at preventing plastic waste from reaching the environment (implemented at post-consumption stages, e.g., waste collection and recycling); and (iii) downstream curative strategies designed to clean legacy pollution in ecosystems where plastic debris has already accumulated (implemented at post-consumption stages, e.g., ocean cleanup). The cost of several strategies belonging to these three categories are presented below. All costs hereinafter are expressed in US\$ at prices for the year 2021 (unless otherwise stated), which explains why the cost data provided in this paper may slightly differ from those in their original publications. Costs estimated over a period of time of several years in this paper are all calculated summing annual costs year-by-year over the period and using a discount rate of 3.5%. Private costs are estimated in Sub-sections 2.1 to 2.3, and external costs and social costs in Section 3 (Table 1 summarizes them). *“The idea underlying the notion of social cost is a very simple one. A man initiating an action does not necessarily bear all the costs (or reap all the benefits) himself. Those that he does bear are private costs; those he does not are external costs. The sum of the two constitutes the social cost”* (de V. Graaf, 2018). Private costs are paid by the firm or the consumer and are included in production and consumption decisions. External costs, on the other hand, are not reflected

on firms' income statements or in consumers' decisions. However, external costs remain costs to society, regardless of who pays for them (Federal Reserve Bank of San Francisco, 2002). Consider a firm or a consumer polluting the marine environment with plastic waste. Because of the firm's or consumer's actions, people regularly eating sea food contaminated with plastics (micro- and nanoplastics) might suffer health effects, tourists may find beaches less attractive due to plastic waste, the beauty of littoral landscapes is damaged, marine animals die through plastic ingestion and entanglement, etc. When external costs like these exist, they must be added to private costs to determine social costs and to ensure that a socially efficient rate of output is generated (i.e., outputs of plastic products and plastic waste).

### *2.1. Upstream solution: stopping plastic production*

A solution that would succeed in reducing plastic emissions into the environment by nearly 100% would consist in entirely stopping plastic production. A report from Grand View Research (2022) estimates the global market share of plastics to be US\$ 593 billion in 2021. Our own calculation (see Section S3 in Supplemental materials) is based on the world input-output table for 2014 (Timmer et al., 2015) and provides results in the same order of magnitude, that is, the global value-added annually produced by the plastic and rubber sector estimated to be US\$ 667 billion in 2021. Hence, if all intermediate consumers (industries and businesses) as well as final consumers (investors, households, public sectors, and non-profit organizations) would stop purchasing plastic products, the global value-added loss would range from US\$ 593 - 667 billion, that is 0.6-0.7% of the world Gross Domestic Product (GDP) in 2021. This is the direct economic cost of stopping plastic production from one day to the next without a transition period. This is a private cost, that is, the cost borne by the producers initiating the action (i.e., shutting down their plastic production activity).

This cost is underestimated since indirect economic costs on suppliers are not considered. Considering them would triple the estimation of the global value-added loss. Indeed, if plastic and rubber production would entirely stop, plastic and rubber industries would have to shut down and their suppliers would no longer be able to sell them energy, raw materials, semi-finished goods and services. Such indirect costs can be taken into account – in addition to direct costs – using Leontief's input-output equations (Leontief, 1936 and 1970; Miller and Blair, 2009, p. 21; Uehara et al., 2018, p. 4). Input-output equations provide further economic details reflecting inter-industrial sales of intermediate inputs between economic sectors (intermediate consumers), in addition to sales to final consumers. We simulated direct and indirect costs of stopping plastic production in the world input-output table (Timmer et al., 2015), which we modified setting to zero the sales of goods and services from plastic and rubber industries to intermediate and final consumers, as well as the purchases of goods and services by plastic and rubber industries from other economic sectors. By using the modified world input-output table to run Leontief's input-output equations (see Supplemental materials, Section S3), we estimate the global GDP loss to be 1.9% in 2021, which includes the direct and indirect costs resulting from entirely stopping plastic and rubber production. This represents an annual loss of US\$ 1875 billion. Such a scenario is unlikely in 2023, as such a drastic solution would require a transition period of several years for the global economic system to adapt to avoid a huge economic cost as well as unavoidable massive employment losses. Plastics are materials used in virtually every sector of manufacturing and use. If plastics production were to cease entirely, there would be a massive disruption in society (which is not taken into account by the Leontief's input-output equations we run), well beyond unemployment and lost sales. However, with the international



United Nations Treaty on Plastic Pollution planned to be finalized in 2024, the political and legislative context might contribute to creating incentives in that direction.

## 2.2. Combining upstream, middle, and downstream solutions: system change scenario

Lau et al. (2020) explain that neither upstream preventive interventions nor downstream curative interventions alone are sufficient to address plastic pollution. Combining the maximum foreseen application of preventive and curative interventions, that is at pre- and post-consumption stages, is the only way to achieve significant plastic pollution reduction in the future (Lau et al., 2020, Cordier et al., 2019). Lau et al. (2020) simulated such a combined scenario, which they named the “*system change scenario*” (SCS). This scenario simulates upstream interventions by considering opportunities to reduce the total plastic quantity produced globally (e.g., through reuse, eliminations such as bans on single-use plastic bags, eliminating plastic overpackaging, etc.) and to substitute plastics with alternative materials (i.e., paper, coated paper and compostable materials). They did not include in the “*system change scenario*” substitute materials that would result in higher life-cycle greenhouse gas emissions compared to plastics (e.g., single-use glass, aluminum and laminated cartons). They also excluded substitute materials with unacceptable health or performance risks (Lau et al., 2020, pp. S18-S22 and Table S20 in their supplemental materials). They assessed the applicability of each reduction and substitution lever to different categories of plastic based on existing businesses, policies, available technologies, environmental trade-offs, and consumer trends observed to date.

Lau et al. (2020) also include mid-stream interventions by simulating improvements to plastic waste collection and disposal systems in order to substantially reduce plastic waste mismanagement (e.g., investments required to replace open dumps by controlled landfills, to increase plastic recycling, etc.). A downstream curative solution is also taken into account in the scenario: beach cleanups to remove plastic debris found in the sand. The full set of their intervention measures is available in Lau et al. (2020, supplemental materials, pp. 71 and 126).

Their results show that annual plastic emissions into the global ecosystem – terrestrial and aquatic together (see Section S1 in Supplemental materials for annual values) – could be decreased by 75-84% in 2040 with the “*system change scenario*” relative to the *business-as-usual* scenario (BAU) (the BAU level is the one that would be achieved if no plastic pollution abatement strategies are undertaken other than those already implemented before 2020). However, when summing annual emissions year-by-year over 1950-2040 to compute accumulated values (using the same calculation method as explained below Figure 2, and Figures S1 and S2 in Supplemental materials), the reduction is much lower. Accumulated emissions of plastic debris over 1950-2040 in the “*system change scenario*” (not shown in Figure 2) amount to 368-574 MMT in aquatic ecosystems and 547-1148 MMT in terrestrial ecosystems, whereas in the BAU scenario they amount to 576-900 MMT and 830-1664 MMT (Figure 2, Lau et al. curves), respectively. This represents a decrease of only 31-36% compared to BAU accumulated levels.

Lau et al. (2020) estimate that from 2016-2040, the total cost of implementing the “*system change scenario*” would be US\$ 470-892 billion (low and high estimate) with a best estimate of US\$ 778 billion. In that scenario, plastic pollution reduction strategies start in 2020 and end in 2040. In the BAU scenario, the total net cost is estimated to be US\$ 953 billion (best estimate) with a low and high estimate of US\$ 643-1077 billion (Lau et al., 2020). The cost estimations in both scenarios cover the cost of collecting, sorting, recycling and disposing of plastic municipal solid waste and are net of revenues

associated with the sale of recycled plastic feedstock and electricity generated from plastic incineration with energy recovery (Lau et al., 2020, p. 9). These estimations are private costs, that is, the cost borne by the municipality (financed by tax payers) or sometimes a private company contracted by the municipality to handle household waste. All these costs are net present value displayed on graphs published in Lau et al. (2020) as well as in their Excel files available in Zenodo (downloadable from this link: <https://zenodo.org/record/3929470>)<sup>1</sup>.

These cost estimates correspond with the level of global discards of inadequately managed plastic waste estimated by Lau et al. (2020)'s model (Figures 1 and S1). However, among all models from Figures 1 and S1, Lau et al. (2020) provides estimates that are among the low and middle curves. Therefore, it might be interesting to consider also high estimates of inadequately managed plastic waste in the estimation of costs in order to reflect the full range of model estimations. If we consider the highest curve in Figure 1, computed based on Lebreton and Andrady (2019), and assuming a direct proportionality between waste management costs and the discard of inadequately managed plastic waste, the net cost of Lau et al.'s scenarios would reach US\$ 643-1612 billion for the BAU scenario ("Inaction scenario" in Table 1) and US\$ 470-1335 billion for the "*system change scenario*" ("Action scenario" in Table 1). This means the "*system change scenario*" is actually US\$ 174-277 billion cheaper than the BAU scenario. In other words, changing the system towards less plastics brings about a benefit, not a cost. This is because although some waste management costs increase in the "system change scenario" compared to the BAU scenario, these additional costs are offset by: (i) revenues from increased quantities of recycled plastic sold by municipalities to the private sector as a raw material (it is usually municipalities that are responsible for collecting and managing household waste) and (ii) savings earned by municipalities from reduced plastic production (because it leads to lower waste production and therefore implies that less waste has to be managed by municipalities, thus reducing plastic waste disposal costs) (Lau et al., 2020, p. 3).

However, other private costs might arise in the private sector, for example involving corporate engagement, through improved product design, alternative material development and new business models that will be necessary to implement the "*system change scenario*" (Lau et al., 2020, p.3). This engagement will require a significant shift in private sector investment through a transition period. The transition cost for the private sector is not estimated in Lau et al. (2020) since their estimate covers only waste management costs, which are generally borne by taxpayers. However, they estimate that in the "*system change scenario*", progressively reducing plastic production and substituting plastics with alternative materials would lead to decreasing plastic production by 47% in 2040. They simulated this scenario assuming a gradual reduction of production through a transition period of 20 years starting in 2021 and ending in 2040. Hence, in the "Action scenario" (Table 1), we reflect that transition period by gradually increasing the reduction by 2.35 percentage points each year compared to the 2021 production level in the BAU scenario. It starts with a reduction percentage of 2.35% in 2021, 4.70% in 2022, 7.05% in 2023, ..., 44.65% in 2039, and 47% in 2040 compared to the BAU production level in 2021. Based on these percentages, we estimated a part of the transition cost for the private sector. If such a production decrease would occur in the plastic industry at the global scale, taking into account the direct effects on plastic industries as well as indirect effects on their suppliers, it would generate a global GDP loss going from 0.05% in 2021 to up to 1.00% in 2040, which represents an annual loss going from US\$ 52.3 billion

---

<sup>1</sup> The Excel files were also sent to us by email in February 2023 by James E. Palardy, one of the authors of Lau et al. (2020)'s article.



in 2021 to US\$ 963.5 billion in 2040. We computed this estimation with the world input-output model mentioned in Section 2.1 (see also Sections S3 and S6 in Supplemental materials). The 20-year transition period allows plastic businesses to take the time required for restructuring and adapting their activity to a low plastic economy. This transition time is also needed for alternative materials markets to grow and replace the vast array of market applications of plastics (offsetting the losses in the traditional plastics industry). Our estimation gives a total cost of transition for the private sector amounting to US\$ 4847-5317 billion (Table 1). This is the total present value estimated with a discount rate of 3.5% over 2021-2040. Some industries will be able to rapidly produce alternative materials and replace plastic materials across the 20-year transition period, which will create positive economic growth opportunities for new businesses. Other businesses will take more time but in any case, annual production of substitute materials are expected to grow every year under the System change scenario from 2.0 million to 62.1 million tons per year across 2021-2040 (low estimate) or from 2.6 million tons/year to 81.1 million tons/year (high estimate) – low and high estimates are provided by Lau et al. (2020) in Zenodo (available here: <https://zenodo.org/record/3929470>). This will generate benefits that are considered in our estimations of the 20-year transition cost. The low and the high estimates of the transition period cost displayed in Table 1 (which are calculated in supplemental materials, Section S6.1) assume that annual production of substitute materials will grow following the low and high estimate ranges provided by Lau et al. (2021), respectively (i.e., 2.0-62.1 million tons/year and 2.6-81.1 million tons/year across 2021-2040, respectively).

298

### 299 2.3. Downstream solution: terrestrial and aquatic ecosystem cleanup

Cost estimations from Lau et al. (2020) presented in Section 2.2 do not include cleanup interventions in aquatic ecosystems. The same for terrestrial ecosystems (only beach cleanups are considered in Lau et al.). However, under the “*system change scenario*”, a large amount of plastic debris still remains in ecosystems due to the legacy pollution. It must be removed if we want damages caused to living organisms (humans included) to stop. Figure 2 shows that the total amount of plastics accumulated at the global scale over 1950-2040 is expected to reach 830-1664 MMT in terrestrial ecosystems (Figure 2 upper graph) and 164-900 MMT in aquatic ecosystems (Figure 2 lower graph) under the BAU scenario. Under the “*system change scenario*”, this amount is expected to drop by 31.0-34.1% in terrestrial ecosystems and by 36.1-36.2% in aquatic ecosystems (Section 2.2). Applying these reduction percentages to the BAU values displayed in Figures 2 and 3 gives an amount of plastic debris accumulated from 1950-2040 under the “*system change scenario*” of 547-1148 MMT for plastic accumulated on terrestrial ecosystems, 0.6-9 MMT for plastics floating in the ocean offshore, 21-331 MMT for plastics floating in the ocean close to the shoreline, 2-37 MMT for plastics sinking to the seabed, 22-56 MMT for plastics floating in rivers, and 49-122 MMT for plastics sinking to lake- and riverbeds. The resulting cleanup cost are calculated in the following paragraphs.

Assuming that beach cleanup practices can be applied to remove plastic debris in all terrestrial ecosystems, we multiply the total amount of plastic accumulated in terrestrial ecosystems under the “*system change scenario*” (calculated in previous paragraph) by the beach cleanup unit cost, which is estimated to be US\$ 1.26-2.06 per kg of plastic collected – unit cost provided by Cruz et al. (2020, p.7) for achieving a degree of cleanliness ranging from clean to very clean. This gives a total present value of US\$ 507-1739 billion (Table 1), which is the private cost to remove the total amount of plastic debris

accumulated over 1950-2040 in terrestrial ecosystems at the global scale under the “*system change scenario*” (starting cleanup activities in 2020 and ending in 2040 as in Lau et al.’s scenario).

Figure 3 (upper graph) shows that plastic debris accumulated in the global ocean will reach 38-590 MMT in 2040 under the BAU scenario. The box on the graph shows that 87.8% of these plastics are floating close to the shoreline and 2.4% are floating offshore. This represents a total amount of 33-518 MMT for plastic debris floating close to the shoreline and of 0.9-14 MMT for plastic debris floating offshore under the BAU scenario. Under the “*system change scenario*”, these amounts are expected to drop to 21-331 MMT for plastic debris floating close to the shoreline and to 0.6-9 MMT for plastic debris floating offshore. The unit cost of the technology developed by The Ocean Cleanup to remove plastics floating offshore is estimated between US\$ 26.6 and US\$ 37.3 per kg of plastic (Tjallema, 2022; The Ocean Cleanup, 2021). The lower margin is the cost The Ocean Cleanup foundation expects to achieve in the short-term based on scaled current technology (System 03), and the higher margin is the cost of the current technology (System 02). To estimate the removal cost of plastics floating offshore, we use this range US\$ 26.6-37.3 per kg. To estimate the removal costs of plastics floating close to the shoreline, we did not find any data. However, we assume this cost to be cheaper than offshore costs since transporting collected plastic debris back to land (to be sent to waste treatment facilities) operates over a much shorter distance than offshore plastics, reducing fuel costs. Therefore, we used the lower unit cost estimated by The Ocean Cleanup foundation, US\$ 16.0 per kg, which is the cost they expect to achieve in the period after optimization (System 04). Based on these unit costs, starting ocean cleanup activities in 2020 and ending in 2040, we estimate US\$ 11-248 billion to be the total present value of the private cost required to remove the total amount of plastic debris floating offshore in the global ocean accumulated over the period of 1950-2040 under the “*system change scenario*”. The total present value of the removal cost for plastics floating close to the shoreline is estimated to be US\$ 251-3895 billion (Table 1).

Here we do not consider the cleanup cost for plastic debris on the seabed (9.8% of plastics accumulated in the ocean – box in Figure 3, upper graph) since the depth and the costs are probably too high to be considered as a serious option. Cleanup of accumulated plastic debris on lake- and riverbeds is not considered either because of lack of robust unit cost data per kg. Figure 3 (lower graph) shows plastic pollution in these environments will reach 76-192 MMT in 2040 under the BAU scenario, twice the amount of plastic floating in rivers (Figure 3, middle graph). This should be considered in a further study.

Figure 3 (middle graph) shows that floating plastic debris accumulated in rivers globally will reach 35-88 MMT in 2040 under the BAU scenario. Under the “*system change scenario*”, this is expected to drop to 22-56 MMT. We multiplied this range by the unit costs of floating plastic removal technologies in rivers (sea bins, trash racks, and booms), which is estimated to be US\$ 1.4-33.3 per kg of plastic removed (Nikiema and Asiedu, 2022, p. 24568). The multiplication gives a total present value of US\$ 23-1373 billion (starting cleanup activities in 2020 and ending in 2040) as the private cost to remove the total amount of floating plastic debris accumulated in rivers from 1950-2040 under the “*system change scenario*”.

All these private costs are summarized in Table 1 and Figure 4 and compared to the cost of inaction, which is estimated in Section 3.

### 3. Global cost of plastic pollution: the cost of inaction

362 Although there is little question about the negative and persistent impacts of plastic pollution on the  
 363 environment (MacLeod et al., 2021), “*how much does it cost*” is a question not well investigated yet. A  
 364 few studies have estimated the global annual cost of plastic pollution in terms of its negative impact on  
 365 the environment. UNEP (2014) was the first to calculate the global cost of plastic pollution, which was  
 366 estimated to be US\$ 89 billion per year. This cost includes plastic-derived environmental damages to  
 367 natural capital through greenhouse gas emissions, water extraction, air, water and land pollution during  
 368 the extraction of natural resources and their conversion into plastic feedstock as well as during plastic  
 369 product end-of-life stages during waste collection and treatment. UNEP (2014) also estimates the  
 370 downstream impact caused by plastic litter leakages into the marine environment, including economic  
 371 losses incurred by fisheries and tourism due to plastic litter (e.g., vessel damage caused by plastic waste  
 372 snarled in a ship’s propellers), loss of amenity caused by litter, time and money spent cleaning up beaches,  
 373 and the ecological cost linked to the loss of species based on monetary valuation approaches, which use  
 374 surveys to estimate how much society would be willing to pay to prevent species loss through plastic  
 375 ingestion and entanglement. They estimate the global cost of plastic litter leakages into marine  
 376 environments to be US\$ 15 billion per year.

377 However, UNEP (2014) calculated these costs before the first estimations of global plastic emissions  
 378 into the ecosystems were provided by scientists, that is, Jambeck et al. (2015), Lebreton et al. (2019),  
 379 Lau et al. (2020), Borrelle et al. (2020) and the OECD (2022). As a result, we decided not to rely on  
 380 UNEP (2014), which recognizes in its report that their cost estimations suffer severe limitations: “*while  
 381 the upstream impacts of producing plastic feedstock are included, the impacts of the manufacturing stage  
 382 are excluded due to their diversity. Downstream impacts, in particular of plastic waste reaching the  
 383 ocean when littered, are likely to be underestimated due to the absence of robust data and scientific  
 384 research [...]*” (UNEP, 2014, pp. 10 and 24).

385 A WWF report authored by de Wit et al. (2021) provides another estimate of the global cost of plastic  
 386 pollution in the marine environment caused by plastic produced in 2019. They estimated this cost to be  
 387 US\$ 2226-4346 billion, with a mid-estimate of US\$ 3286 billion. However, as explained by de Wit et al.  
 388 (2021, p. 38), the WWF report’s estimation relies on and extrapolates from a scientific article published  
 389 by Beaumont et al. (2019). Thereby, we decided to rely directly and exclusively on Beaumont et al.  
 390 (2019) in our paper.

391 Beaumont et al. (2019) estimated the global annual cost of plastic pollution in the marine environment  
 392 to be US\$ 3975-39753 per ton of marine plastic. (This global cost slightly differs from the original data  
 393 provided in Beaumont et al. (2019) because, as mentioned in Section 2, all costs in our paper are  
 394 expressed in US\$ at prices for the year 2021 unless otherwise stated). Their estimations are external costs  
 395 (see definition in first paragraph of Section 2) related to non-market ecosystem services. They exclusively  
 396 considered the depreciation of marine natural capital - marine ecosystem services - caused by plastic  
 397 pollution. The estimation from Beaumont et al. (2019) relies on a semi-systematic literature review of  
 398 1191 data points, which they used to compute the impact scores of plastic pollution on marine ecosystem  
 399 services by subject type (e.g., turtles, birds, fish, etc.). The ecosystem services they considered cover  
 400 three categories: provisioning, regulatory and cultural services following CICES’s classification (CICES,  
 401 2013). However, the fourth category, supporting services (Millennium Ecosystem Assessment, 2005), is  
 402 lacking in Beaumont et al.’s estimation. The impact scores were translated into monetary values in 2011  
 403 by using the global database for ecosystem services values based on benefit transfer techniques (Costanza  
 404 et al., 2014). Benefit transfer is a well-known monetary valuation technique used in environmental  
 405 economics to estimate the economic value of ecosystem services for which no money is exchanged on a

market (Pearce et al., 2006). For comparison with plastic reduction strategies estimated in Section 2, we multiplied year-by-year the total amount of plastic debris accumulated in the ocean with the global annual cost per ton of marine plastics across 25 years over the 2016-2040 period (using a discount rate of 3.5%, as for all other costs calculated over a period of time of several years in this paper). In the multiplication, for the amount of plastic debris accumulated over years, we used the highest estimation from Lebreton et al. (2019) and the lowest one from the OECD (2022) (Fig. 3, upper graph). It gives a total global cost over the 2016-2040 period ranging from US\$ 1862 billion to US\$ 268498 billion for the “Inaction scenario” and from US\$ 1003 billion to US\$ 132819 billion for the “Action scenario” (Table 1). The “Action scenario” causes damages to the ecosystems too (although its environmental cost is reduced by half compared to the “Inaction scenario”) because preventive and clean-up operations described in Section 2 take time. They are implemented progressively on an annual basis. Meanwhile although plastic pollution is gradually reduced, plastic debris approaches the zero level in ecosystems by 2040 (see Figure S4 in Supplemental materials). And since plastic sinking on sea-, lake- and riverbeds are not cleaned up in the “Action scenario”, a residual amount remains present in the ecosystems by 2040 (between 3 and 36 MMT in the global ocean in 2040 under the “Action scenario” – Figure S4 in supplemental materials). Moreover, the “Action scenario” strongly reduces annual emissions of plastic debris to ecosystems (by 75 to 84 % compared to BAU scenario levels, see Section 2.2) but it does not completely stop them. The “tap” of plastic pollution is not completely turned off.

Plastics also have important effects on public health due to endocrine-disrupting chemicals found as additives in plastic products, which are suspected to cause several diseases: IQ loss and intellectual disability, adult diabetes, endometriosis, obesity, cryptorchidism (undescended testicle in the scrotum), male infertility, low birth weight, pneumonia, kidney cancer, hypothyroidism, polycystic ovarian syndrome, breast cancer, and low testosterone resulting in increased early mortality. Diseases due to chemicals used in plastic materials is substantial, costing US\$ 384-403 billion each year in the USA, US\$ 44 billion per year in the European Union (United Kingdom included), and US\$ 18 billion per year in Canada. These three estimates are external costs for diseases that occurred in 2010 and are expressed in US\$ at the price of the year 2010 (see more information in Supplemental materials in Section S5 compiled by the Endocrine Society based on Trasande et al., 2015, 2016, 2022, 2022a; Gore et al., 2015; Attina et al., 2016; Malits et al., 2022; Obsekovic et al., 2022). Converted into US\$ at 2021 prices<sup>2</sup> and summed across USA, EU and Canada, gives a total annual cost of US\$ 553-577 billion. Assuming this total annual cost is constant and summed year-by-year over 2016-2040 gives a total present value of that cost as US\$ 11206-11692 billion. This estimation is conservative given the annual cost is likely not constant. Population growth and plastic production growth probably will lead to increases in the annual number of people affected by plastic-related diseases and annual public health costs. In addition, such cost estimations should be carried out for all regions of the world to obtain a global human health cost. Due to the lack of studies, we had to neglect the rest of the world and consider only the USA, the EU (United Kingdom included) and Canada. In the report from UNEP (2023), Landrigan et al. (2023, p. 100) provide other estimates of public health costs related to plastic additives. Most of them are based on the same publications as those we use from the Endocrine Society. For this reason, we decided to rely directly and exclusively on Endocrine Society data (in supplemental materials, Section S5). Landrigan et al. (2023, pp. 99 and 102), UNEP (2023, p. 6) and Merkl and Charles's (2022) estimated other health costs related to plastics: the economic costs of deaths of workers attributable to ambient particulate matter air pollution

---

<sup>2</sup> Conversion rate for inflation: 1 US \$ in 2010 = 1,24 US\$ in 2021.



(PM<sub>2.5</sub>) and to occupational exposure resulting from plastics production. Merkl and Charles (2022) also estimated the social cost of carbon emitted during plastic production. These estimations are not taken into account in our paper but could be considered in further research.

The total health cost estimation mentioned in the previous paragraph (US\$ 11206-11692 billion) is taken into account in the “Inaction scenario” and the “Action scenario” as well (Table 1). We made this choice because in the “Action scenario”, plastic pollutants do not tend to zero before 2040. As explained above, plastic (pollution and production) reduction strategies are implemented progressively on a year-by-year basis. Thus, people are continuously exposed to plastics, although to a diminishing extent, across the period 2016-2040. In addition, diseases due to exposure to plastics are not caused only by pollutants but also by plastics products (especially food packaging and plastic bottles) to which humans are frequently exposed. And yet, in the “Action scenario”, these plastic products, although their production is reduced by almost half, they are not entirely eliminated. A full epidemiologic-economic study would be required to estimate the potential reduction in human exposure in the “Action scenario” and the effect on health cost. Therefore, this has not been taken into account, which explains why the health cost in the “Action scenario” is probably overestimated.

The last cost we include in the calculation of the global cost of inaction comes from Lau et al. (2020). As mentioned in Section 2.2, they estimated waste management costs in the case of inaction between US\$ 643-1612 billion, which is greater than in the case of action.

Summing these three categories of costs (marine pollution, public health, and waste management) gives a total global cost over the 2016-2040 period ranging from US\$ 13711 billion to US\$ 281802 billion, that is US\$ 548-11272 billion per year when divided by the 25 years of the period. This annual range is wider than the one presented in UNEP (2023, pp. 6 and 8), which is estimated to be US\$ 294-1500 billion per year. The first reason is because we directly use the unit cost of damages (cost per tons of plastic debris) caused to ecosystems estimated by Beaumont et al. (2019) whereas UNEP (2023, pp. 6 and 8) uses the unit cost from WWF (de Wit et al., 2021). The WWF study estimated the impact of marine plastic debris caused by plastic produced in 2019, whereas we estimate the impacts caused all years across the 2016-2040 period due to plastic debris accumulated in marine ecosystems since 1950. In addition, for the calculation of the total cost, UNEP (2023, pp. 6 and 8) multiplied the unit costs by the amount of plastic pollution estimated by the Pew Charitable Trusts and Systemiq (2020), which is the report version of the scientific article published in Science by Lau et al. (2020). In our paper, we base our calculations on a set of eight global plastic models estimating plastic pollution (Figure 1, 2 and 3): Jambeck et al. (2015), Lebreton and Andrady (2019), Cordier and Uehara (2019), Lau et al. (2020), Borelle et al. (2020) Cordier et al. (2021), OECD (2022), and Yan et al. (2022).

#### 4. Global benefits obtained from plastics

In this Section, we compare the costs calculated in Sections 2 and 3 (Summarized in Table 1) to the benefits obtained from plastics in the form of income, that is, wages and salaries for workers, dividends for investors, rents for building owners, taxes for government budgets, etc. In Section 2.1, we calculated the global direct and indirect contribution of the plastic industry on global GDP, which we estimated to be US\$ 1875 billion in 2021. This represents the annual benefit plastic products bring about as income to individuals involved in economic activities linked to plastics. Summing this annual benefit across the period 2016-2040 gives a total of US\$ 37985 billion in the case of inaction. Subtracting from this



estimation the total cost of transition for the private sector, that is, US\$ 4847-5317 billion (calculated with the world input-output model mentioned in Section 2.2, last paragraph), yields US\$ 32668-33138 billion, which is the benefit earned in the case of actions towards zero plastic pollution by 2040. This represents a 13-14% loss compared to the “Inaction scenario”. These amounts are summarized in the three first columns of Table 2.

Benefits can be converted into net benefits by subtracting the costs (costs calculated in Section 3 and Table 1) from the benefits (first three columns in Table 2). We made this calculation for the “Action” and the “Inaction” scenarios (using the costs calculated in Section 2.2-2.3 and 3, respectively). This yields the two last columns in Table 2 and shows that in the case of action towards zero plastics by 2040 (including 47% reduction of plastic production by 2040), net benefits might be either negative or positive, ranging from US\$ –120433 billion to US\$ 19667 billion. The positive estimate means action towards zero plastic pollution is a gain for the global community altogether (private sectors, public sector, civil society, and ecosystems). The negative estimate represents a cost for the global community. In the case of inaction, we face a similar situation: the net benefit might be either positive or negative and is expected to be between US\$ –243817 billion and US\$ 24274 billion. The high estimate, that is the positive net benefit, means that inaction might bring about benefits that offset the global environmental costs generated by plastic pollution in case of inaction. The low estimate indicates negative net benefit, that is to say, the dramatic costs that may be incurred through inaction.

## 5. Discussion and conclusion

### 5.1. Comparison of the cost of action and inaction

Table 1 summarizes the costs that will be incurred if the plastic pollution intervention strategies presented in Section 2 are implemented between 2020 and 2040. It also displays in the penultimate row the cost of global plastic pollution estimated in Section 3 under the BAU scenario. Table 1 and Figure 4 show the global cost of a combination of actions towards zero plastic pollution undertaken in all countries by 2040 to be US\$ 18.3-158.4 trillion (which includes reducing plastic production by 47% in 2040, replacing plastic products with alternative materials, improving waste collection and treatment, and cleaning up ecosystems). If no actions are undertaken, the cost of damages caused by plastic pollution from 2016-2040 is estimated to be US\$ 13.7-281.8 trillion. This suggests inaction could generate a global cost either 1.3 times cheaper than the cost of action or up to 1.8 times more expensive.

Plastic product sales will also generate a global benefit in the form of incomes (salaries, dividends, taxes, etc.) estimated to be US\$ 37.99 trillion from 2016-2040 in case of inaction and US\$ 32.67-33.14 trillion in the case of action. Calculating benefit minus costs provides net benefits of US\$ –120.43-19.67 trillion in the case of action and US\$ –243.82-24.27 trillion in the case of inaction (Table 2 and Figure 5). This suggests action and inaction will be beneficial only considering the high estimate. The low estimates are both negative (US\$ –120.43 trillion and US\$ –243.82 trillion for action and inaction, respectively), which means action and inaction might generate a net cost for the entire society. In the case of inaction, it is because benefits obtained from plastic products will not be sufficient to offset costs of plastic pollution impacts; in the case of action, it is because reduced ecosystem damage costs will not be sufficient to offset the cost of actions towards zero plastic pollution.

However, the global damage cost estimated in our paper (penultimate row of Table 1) is significantly underestimated. We therefore cannot exclude the possibility that future studies will show a negative value for the higher estimate of the net benefit in the case of inaction (meaning that it would be a net cost and not a net benefit).

Three reasons explain the underestimate of the cost of global environmental damages in case of inaction. First, the estimated cost of global damages caused by plastics exclusively covers marine ecosystems and omits terrestrial ecosystems. There is an urgent need to develop studies on the cost of plastic contamination on land. The cost of global damage caused by plastic pollution to terrestrial ecosystems is likely to be significant given the total amount of plastic debris that will accumulate on land over the 1950-2040 period (830-1664 MMT, Figure 2 upper graph) is higher than in marine ecosystems (38-590 MMT, Figure 3 upper graph).

Second, the cost of plastics on human health is strongly underestimated in our paper since we had to limit the estimation to three countries for which data were available: the USA, the European Union, and Canada. Extrapolating to the rest of the world proportionally to population size is not possible, not even for a restricted set of similar countries such as high-income countries. As underlined by Leonardo Trasande (personal communication by email, 6<sup>th</sup> of June 2023), country-level exposures to plastic additives vary widely by policy context, which explains why the number of people suffering diseases and health costs related to plastic additives are significantly different from one country to another, even within high-income countries.

Third, because of lacking data, except for the model results from Lau et al. (2020), the models displayed in Figures 2 and 3 (and Figures S2 and S3 in the Supplemental materials) do not consider emissions of primary microplastics into the environment (e.g., synthetic textile fibers from washing machines). Further studies should quantify primary microplastic emission to the ecosystem since they are likely to be significant. For example, primary microplastic leakages from tire wear may contribute 5–10% of global ocean plastics loading (Kole et al., 2017; Hale et al., 2020). And even if we could count them, technologies to clean up micro- and nanoplastics in ecosystems are lacking anyway. This explains why we could not estimate the cost of cleaning up these small pieces of plastic debris to remove them from contaminated ecosystems.

The global cost of private sector action estimated in Table 1 also suffers from inaccuracies under the “Action scenario”. First, we estimated with an input-output model the transition cost for the private sector adapting to a low plastic society (Section 2.2, and row 9 in Table 1). The issue is that the input-output model we used is static and assume fixed prices and technology. This does not allow for flexibility in the input-output table, which cannot reflect the way the global economic structure will change due to future technological developments of substitutes and substitute approaches to meeting the decreasing demand for plastics over the coming decades under the « Action scenario ». This likely means the transition costs are over-estimated. We must, thereby, acknowledge the limitations of using static input-output models for benefit-cost analyses over multi-decadal timelines (beyond a 10-year period, the technological changes are likely to be significant, which is hardly captured by static input-output models). Further research could solve this drawback by dynamising input-output technological coefficients (e.g., Uehara et al., 2018) or, as recommended by the U.S. Environmental Protection Agency (US EPA 2014 and 2020, pages 8-9 to 8-21 and 8-16 to 8-26, respectively), by using a computable general equilibrium model. Second, we estimated the economic impacts of ocean, river and terrestrial cleanups (on the cost side) based on operational cost of removing plastic debris from the ecosystems (Section 2.3, and rows 3-6 in Table 1). However, these are the direct costs. Indirect costs have not been taken into consideration, since cleanup costs were not passed through the input-output model to reflect the impact on suppliers,

intermediate and final consumers, wages and salaries, etc. Thereby, cleanup costs are likely understated. These two inaccuracies (on transition and cleanup costs) affect the estimates of benefits in Table 2, which consist in calculating differences from the input-output model.

## 5.2. Cost distribution across countries from the global south and global north

The global costs displayed in Table 1 and Figure 4 will not be evenly distributed between Global South and Global North countries. First, as global plastics production continues increasing, this growth is unequally distributed. From 2009 to 2019, annual global plastics production grew from 321 MMT to 460 MMT (OECD, 2022, p.68). During the same period in Europe, production was comparably stable, increasing from 55 MT in 2009 to 58 MMT in 2019 (PlasticsEurope, 2011 and 2020) in response to increasing social and environmental regulation.

Second, plastic waste management also reflects planetary asymmetry in how the benefits and harms of plastics are distributed. For example, prior to 2018, China imported over half of the world's plastic waste. In 2018, when China began implementing their near total ban on plastic waste imports, the resulting reshuffling of the global plastic waste market resulted in other countries, including some of the world's poorest, such as Malaysia, Thailand, and Indonesia, importing much larger quantities of global plastic waste (Vidal, 2020), and the associated consequences for ecosystems and human health in these countries (Marrs et al., 2019; Trasande, 2022). An estimated 58% of all plastic produced between 1950 and 2017 has been discarded and continues impacting the environment (Geyer, 2020). As plastic production continues increasing, so too do the negative impacts of plastic-derived pollution throughout plastics' material life cycle. While some countries are introducing plastic-related regulation, so long as plastic production continues increasing, the harmful socioecological consequences of plastics will be displaced to less-regulated countries, such as Turkey or Romania in the case of Europe, and Malaysia, Thailand and Indonesia in Asia, not to mention the globally shared consequences for the Earth's oceans and climate.

Third, the economic impact of the cost of future mitigation policies will probably be uneven across countries as a recent study by the OECD (2022) shows. The study considers a wide range of policies intended to restrain plastic production and consumption as well as to enhance design for circularity (e.g., plastic tax, eco-design for durability and repair), improve recycling (e.g., recycled content targets), and close leakage pathways (e.g., better plastic waste collection). The degree of effort varies by country's income level. The "*global ambition policy scenario*" simulated by the OECD (2022) intends to reduce plastic leakage to the environment to nearly zero by 2060. The costs resulting from this scenario incurred by Global South countries (red bars in Figure 6) will be among the highest (except in China). For example, in Sub-Saharan African countries, GDP is projected to decline by 2.8 % below the baseline. The Global North (blue bars in Figure 6) will be much less affected. For example, in OECD EU countries (that is, high-income countries), GDP is projected to decline by only 0.2% below the baseline, mostly because the economic infrastructure in OECD countries, waste collection and treatment infrastructures included, is fundamentally more extensive than in non-OECD countries). Non-OECD EU countries (labeled "other EU" in Figure 6) are, however an exception in the Global North since their GDP is projected to decline by 2.1% below the baseline. One of the reasons for sharp GDP declines in Global South and non-OECD EU countries is due to substantial investments still missing that are required to improve waste collection and treatment facilities to achieve the policy targets set in the scenario.

## 5.3. What to do now?

624 Knowing that any plastic production implies pollution in different forms across different scales, and that  
 625 the producers' intentions are to increase their own benefits, as demonstrated by the past 50 years of  
 626 production:

- 627     – Why would producers agree to reduce their otherwise growing benefits?  
 628         How will such public policy get implemented against private sector interests? As things stand  
 629         currently, it is not easy to do so.
- 630     – The macroeconomic models and global estimates create abstractions far from local, regional, and  
 631         national realities, proposing dialogue/s between developed and developing countries as  
 632         solution/s, when today the questions can be focused more on: "Where do the benefits go?" "Where  
 633         are the impacts?" and "Who has the capacity to regulate the asymmetry?"

634 In some regions, producers and recyclers are the same corporate entity, giving them an interest in  
 635 maintaining growth from both sides (production and recycling). We do not see how they can then be part  
 636 of the solution on their own. Corporations have committed documented abuses for decades, everything  
 637 from greenwashing to murder, and continue doing so today. This is well documented for longer-running  
 638 environmental concerns such as climate change, mining, or asbestos (Bonneuil et al., 2021; Supran et al.,  
 639 2023; Middeldorp and Le Billon, 2019; Le Billon and Lujala P., 2020; Forbidden stories, 2019; Ladou,  
 640 2004). Similar publications on the role of plastic-related corporations (e.g., soft drink industries) are still  
 641 in their infancy (Wood et al., 2021; Dauvergne et al., 2018). However, it is progressing since the scandal  
 642 of the leaked internal document from Coca-Cola (Coca-Cola Europe, 2016) revealing the company  
 643 prioritized a "fight back" strategy against EU policies that planned to implement EPR schemes (Extended  
 644 Producer Responsibility), to increase plastic collection and recycling, and to develop deposit return  
 645 schemes. We must avoid denial about this, keeping in mind a sentence from the trials against the tobacco  
 646 industry in the 1990s when the U.S. District Judge H. Lee Sarokin said in 1992: "Who are these persons  
 647 who knowingly and secretly decide to put the buying public at risk solely for the purpose of making  
 648 profits, and who believe that illness and death of consumers is an appropriate cost of their own  
 649 prosperity!" (Brownell and Warner, 2009).

650 It comes down to this: allowing plastic production, consumption, or recycling to continue growing means  
 651 allowing plastic pollution and its associated costs to continue increasing (Trasande, 2022). While there  
 652 are gaps in the data, the estimates provided here illustrate the high economic costs of inaction regarding  
 653 plastic pollution, along with the need to ensure the costs of addressing plastic pollution are not inequitably  
 654 born by those least responsible, who have benefited least.

655

## 656 **Acknowledgements**

657 We are grateful to the anonymous reviewers who helped to considerably improve this paper.

## 658 **Author contribution statement:**

659 **Mateo Cordier:** Conceptualization, Data collection, Formal analysis, Methodology, Supervision,  
 660 Validation, Visualization, Writing – original draft, Writing – review & editing. **Takuro Uehara:** Funding  
 661 acquisition, Project administration, Validation, Writing – original draft, Writing – review & editing.  
 662 **Bethany Jorgensen:** Validation, Writing – original draft, Writing – review & editing. **Juan Baztan:**  
 663 Context, writing and review.

664 **Financial support:** This research received support from the JSPS KAKENHI (Japan Society for the  
665 Promotion of Science), grant number 19KK0271.

666 **Competing interests:** The authors declare no competing interests.

667

## 668 References

- 669 Attina TM, Hauser R, Sathyanarayana S, Hunt PA, Bourguignon JP, Myers JP, ... and Trasande L  
670 (2016) Exposure to endocrine disrupting chemicals in the USA: a population-based disease burden and  
671 cost analysis. *Lancet Diabetes Endocrinol* **4**, 996–1003. [https://doi.org/10.1016/S2213-8587\(16\)30275-](https://doi.org/10.1016/S2213-8587(16)30275-3)  
672 [3](https://doi.org/10.1016/S2213-8587(16)30275-3)
- 673 Baekland LH (1909) The synthesis, constitution, and uses of bakelite. *Journal of Industrial and*  
674 *Engineering Chemistry* **1**(3), 149-161. <https://doi.org/10.1021/ie50003a004>
- 675 Bajt O (2021) From plastics to microplastics and organisms. *FEBS Open bio* **11**(4), 954-966. doi:  
676 10.1002/2211-5463.13120
- 677 Beaumont NJ, Aanesen M, Austen MC, Börger T, Clark JR, Cole M., ... and Wyles KJ (2019) Global  
678 ecological, social and economic impacts of marine plastic. *Marine pollution bulletin* **142**, 189-195.  
679 <https://doi.org/10.1016/j.marpolbul.2019.03.022>
- 680 Bonneuil C, Choquet PL and Franta B (2021) Early warnings and emerging accountability: Total's  
681 responses to global warming, 1971–2021. *Global Environmental Change* **71**, 102386.  
682 <https://doi.org/10.1016/j.gloenvcha.2021.102386>
- 683 Borrelle SB, Ringma J, Law KL, Monnahan CC, Lebreton L, McGivern A, ... and Rochman CM  
684 (2020) Predicted growth in plastic waste exceeds efforts to mitigate plastic pollution. *Science*  
685 **369**(6510), 1515-1518. <https://doi.org/10.1126/science.aba3656>
- 686 Brownell KD and Warner KE (2009) The perils of ignoring history: Big Tobacco played dirty and  
687 millions died. How similar is Big Food? *The Milbank Quarterly* **87**(1), 259-294. doi: [10.1111/j.1468-](https://doi.org/10.1111/j.1468-0009.2009.00555.x)  
688 [0009.2009.00555.x](https://doi.org/10.1111/j.1468-0009.2009.00555.x)
- 689 Chamas A, Moon H, Zheng J, Qiu Y, Tabassum T, Jang JH, ... and Suh S (2020) Degradation rates of  
690 plastics in the environment. *ACS Sustainable Chemistry & Engineering* **8**(9), 3494-3511.  
691 <https://doi.org/10.1021/acssuschemeng.9b06635>
- 692 Chen X and Yan N (2020) A brief overview of renewable plastics. *Materials Today Sustainability* **7-8**,  
693 100031. <https://doi.org/10.1016/j.mtsust.2019.100031>
- 694 Chiba S, Saito H, Fletcher R, Yogi T, Kayo M, Miyagi S, ... and Fujikura K (2018) Human footprint in  
695 the abyss: 30 year records of deep-sea plastic debris. *Marine Policy* **96**, 204-212.  
696 <https://doi.org/10.1016/j.marpol.2018.03.022>
- 697 Choy CA, Robison BH, Gagne TO, Erwin B, Firl E, Halden RU, ... and Van Houtan KS (2019) The  
698 vertical distribution and biological transport of marine microplastics across the epipelagic and  
699 mesopelagic water column. *Scientific reports* **9**(1), 7843. <https://doi.org/10.1038/s41598-019-44117-2>
- 700 CICES (2013) Common International Classification of Ecosystem Services (CICES). Retrieved from  
701 [www.cices.eu](http://www.cices.eu)
- 702 Coca-Cola Europe (2016) Radar screen of EU public policies. Monthly issue update: February &  
703 March 2016. 15 pp. Retrieved from: [https://unearthed.greenpeace.org/2017/01/25/investigation-coca-](https://unearthed.greenpeace.org/2017/01/25/investigation-coca-cola-fight-back-plans-tackle-plastic-waste/)  
704 [cola-fight-back-plans-tackle-plastic-waste/](https://unearthed.greenpeace.org/2017/01/25/investigation-coca-cola-fight-back-plans-tackle-plastic-waste/)



- 705 Costanza R., de Groot R, Sutton P, van der Ploeg S, Anderson SJ, Kubiszewski I, Farber S and Turner  
 706 KR (2014) Changes in the global value of ecosystem services. *Global Environmental Change* **26**, 152–  
 707 158. <https://doi.org/10.1016/j.gloenvcha.2014.04.002>
- 708 Cordier M and Uehara T (2019) How much innovation is needed to protect the ocean from plastic  
 709 contamination? *Science of the total environment* **670**, 789–799.  
 710 <https://doi.org/10.1016/j.scitotenv.2019.03.258>
- 711 Cordier M, Uehara T, Baztan J, Jorgensen B and Yan H (2021) Plastic pollution and economic growth:  
 712 The influence of corruption and lack of education. *Ecological economics* **182**, 106930.  
 713 <https://doi.org/10.1016/j.ecolecon.2020.106930>
- 714 Cruz CJ, Muñoz-Perez JJ, Carrasco-Braganza MI, Pouillet P, Lopez-Garcia P, Contreras A and Silva R  
 715 (2020). Beach cleaning costs. *Ocean & Coastal Management* **188**, 105118.  
 716 <https://doi.org/10.1016/j.ocecoaman.2020.105118>
- 717 de Wit W, Burns ET, Guinchard JC and Ahmed N (2021) Plastics: the costs to society, the environment  
 718 and the economy. Gland (Switzerland): World Wide Fund for Nature (WWF).
- 719 Dauvergne P (2018) Why is the global governance of plastic failing the oceans? *Global Environmental*  
 720 *Change* **51**, 22-31. <https://doi.org/10.1016/j.gloenvcha.2018.05.002>
- 721 Forbidden Stories (2019) Green Blood. Video documentary by investigative journalists. Available at:  
 722 <https://forbiddenstories.org/case/green-blood/>
- 723 de V. Graaff J (2018) Social Cost. In: The New Palgrave Dictionary of Economics. London : Palgrave  
 724 Macmillan, pp. 12516-12520. [https://doi.org/10.1057/978-1-349-95189-5\\_1459](https://doi.org/10.1057/978-1-349-95189-5_1459)
- 725 Federal Reserve Bank of San Francisco (2002) What is the difference between private and social costs,  
 726 and how do they relate to pollution and production? Available at:  
 727 [https://www.frbsf.org/education/publications/doctor-econ/2002/november/private-social-costs-](https://www.frbsf.org/education/publications/doctor-econ/2002/november/private-social-costs-pollution-production/)  
 728 [pollution-production/](https://www.frbsf.org/education/publications/doctor-econ/2002/november/private-social-costs-pollution-production/)
- 729 Geyer R, Jambeck JR, Law KL (2017) Production, use, and fate of all plastics ever made. *Science*  
 730 *Advances* **3** (7), e1700782. DOI: [10.1126/sciadv.1700782](https://doi.org/10.1126/sciadv.1700782)
- 731 Geyer R (2020) Production, use, and fate of synthetic polymers. In: Plastic waste and recycling.  
 732 Academic Press, pp. 13-32. <https://doi.org/10.1016/B978-0-12-817880-5.00002-5>
- 733 Gore AC, Chappell VA, Fenton SE, Flaws JA, Nadal A, Prins GS, Toppari J and Zoeller RT (2015)  
 734 EDC-2: The Endocrine Society's Second Scientific Statement on Endocrine-Disrupting Chemicals.  
 735 *Endocrine Reviews* **36**(6), 593-602. DOI: [10.1210/er.2015-1093](https://doi.org/10.1210/er.2015-1093)
- 736 Grand View Research (2022) Plastic Market Size, Share & Trends Analysis Report By Product (PE,  
 737 PP, PU, PVC, PET, Polystyrene, ABS, PBT, PPO, Epoxy Polymers, LCP, PC, Polyamide), By  
 738 Application, By End Use, And Segment Forecasts, 2022 – 2030. San Francisco: Bulk chemicals.  
 739 Available at: [https://www.grandviewresearch.com/industry-analysis/global-plastics-](https://www.grandviewresearch.com/industry-analysis/global-plastics-market/segmentation)  
 740 [market/segmentation](https://www.grandviewresearch.com/industry-analysis/global-plastics-market/segmentation)
- 741 Hale RC, Seeley ME, La Guardia MJ, Mai L and Zeng EY (2020) A global perspective on  
 742 microplastics. *Journal of Geophysical Research: Oceans* **125**(1), e2018JC014719.  
 743 <https://doi.org/10.1029/2018JC014719>
- 744 Jambeck JR, Geyer R, Wilcox C, Siegler TR, Perryman M, Andrady A, ... and Law KL (2015) Plastic  
 745 waste inputs from land into the ocean. *Science* **347**(6223), 768-771.  
 746 <https://doi.org/10.1126/science.1260352>

- 747 Kole PJ, Lohr AJ, van Belleghem F and Ragas AMJ (2017) Wear and tear of tyres: A stealthy source of  
 748 microplastics in the environment. *International Journal of Environmental Research and Public Health*  
 749 **14**(10). <http://doi.org/10.3390/ijerph14101265>
- 750 Lacerda ALDF, Rodrigues LDS, Van Sebillle E, Rodrigues FL, Ribeiro L, Secchi ER, ... and Proietti  
 751 MC (2019) Plastics in sea surface waters around the Antarctic Peninsula. *Scientific reports* **9**(1), 3977.  
 752 <https://doi.org/10.1038/s41598-019-40311-4>
- 753 LaDou J (2004) The asbestos cancer epidemic. *Environmental health perspectives* **112**(3), 285-290.  
 754 <https://doi.org/10.1289/ehp.6704>
- 755 Lau WW, Shiran Y, Bailey RM, Cook E, Stuchtey MR, Koskella J, ... and Palardy JE (2020).  
 756 Evaluating scenarios toward zero plastic pollution. *Science* **369**(6510), 1455-1461.  
 757 <https://doi.org/10.1126/science.aba9475>
- 758 Landrigan PJ, Raps H, Cropper M, Bald C, Brunner M, Canonizado EM, ... and Dunlop S (2023) The  
 759 Minderoo-Monaco Commission on Plastics and Human Health. *Annals of Global Health* **89**(1), 1–215.  
 760 <https://doi.org/10.5334/aogh.4056>
- 761 Law KL and Narayan R (2022) Reducing environmental plastic pollution by designing polymer  
 762 materials for managed end-of-life. *Nature Reviews Materials* **7**(2), 104-116.  
 763 <https://doi.org/10.1038/s41578-021-00382-0>
- 764 Le Billon P, Lujala P (2020) Environmental and land defenders: Global patterns and determinants of  
 765 repression. *Global Environmental Change* **65**, 102163.  
 766 <https://doi.org/10.1016/j.gloenvcha.2020.102163>
- 767 Lebreton L and Andrady A (2019) Future scenarios of global plastic waste generation and disposal.  
 768 *Palgrave Communications* **5**(1), 1-11. <https://doi.org/10.1057/s41599-018-0212-7>
- 769 Lebreton L, Egger M and Slat B (2019) A global mass budget for positively buoyant macroplastic  
 770 debris in the ocean. *Scientific reports* **9**(1), 1-10. <https://doi.org/10.1038/s41598-019-49413-5>
- 771 Leontief WW (1936) Quantitative input and output relations in the economic system of the United  
 772 States. *The Review of Economic Statistics* **18** (3), 105–125. <https://doi.org/10.2307/1927837>
- 773 Leontief WW (1970) Environmental repercussions and the economic structure: an input–output  
 774 approach. *Review of Economics and Statistics* **52**(3), 262–271. <https://doi.org/10.2307/1926294>
- 775 Leslie HA, Van Velzen MJ, Brandsma SH, Vethaak AD, Garcia-Vallejo JJ and Lamoree MH (2022)  
 776 Discovery and quantification of plastic particle pollution in human blood. *Environment international*  
 777 **163**, 107199. <https://doi.org/10.1016/j.envint.2022.107199>
- 778 MacLeod M, Arp HPH, Tekman MB and Jahnke A (2021) The global threat from plastic pollution.  
 779 *Science* **373**(6550), 61-65. <https://doi.org/10.1126/science.abg5433>
- 780 Malits J, Naidu M, Trasande L (2022) Exposure to Endocrine Disrupting Chemicals in Canada:  
 781 Population-Based Estimates of Disease Burden and Economic Costs. *Toxics* **10**(3), 146.  
 782 <https://doi.org/10.3390/toxics10030146>
- 783 Marrs DG, Ručevska I, Villarrubia-Gómez P (2019) Controlling Transboundary Trade in Plastic waste.  
 784 GRID-Arendal. Available at: <https://grid.cld.bz/Controlling-Transboundary-Trade-in-Plastic-Waste>
- 785 Merkl A and Charles D (2022) The Price of Plastic Pollution: Social Costs and Corporate Liabilities.  
 786 Minderoo Foundation. Available at: [https://www.unepfi.org/wordpress/wp-](https://www.unepfi.org/wordpress/wp-content/uploads/2022/10/The-Price-of-Plastic-Pollution.pdf)  
 787 [content/uploads/2022/10/The-Price-of-Plastic-Pollution.pdf](https://www.unepfi.org/wordpress/wp-content/uploads/2022/10/The-Price-of-Plastic-Pollution.pdf)

- 788 Middeldorp N, Le Billon P (2019) Deadly environmental governance: authoritarianism, eco-populism,  
789 and the repression of environmental and land defenders. *Annals of the American Association of*  
790 *Geographers* **109**(2), 324-337. <https://doi.org/10.1080/24694452.2018.1530586>
- 791 Millennium Ecosystem Assessment (2005) Ecosystems and Human Well-being: Synthesis. Washington  
792 DC: Island Press. Available at:  
793 <https://www.millenniumassessment.org/documents/document.356.aspx.pdf>
- 794 Miller RE and Blair PD (2009) Input-output analysis: foundations and extensions. New-York:  
795 Cambridge university press. Available at: [https://www.cambridge.org/core/books/inputoutput-](https://www.cambridge.org/core/books/inputoutput-analysis/69827DA658E766CD1E17B1A47BA2B9C3)  
796 [analysis/69827DA658E766CD1E17B1A47BA2B9C3](https://www.cambridge.org/core/books/inputoutput-analysis/69827DA658E766CD1E17B1A47BA2B9C3)
- 797 Nikiema J and Asiedu Z (2022) A review of the cost and effectiveness of solutions to address plastic  
798 pollution. *Environmental Science and Pollution Research* **29**(17), 24547-24573.  
799 <https://doi.org/10.1007/s11356-021-18038-5>
- 800 Obsekov V, Kahn LG, Trasande L. (2022) Leveraging Systematic Reviews to Explore Disease Burden  
801 and Costs of Per- and Polyfluoroalkyl Substance Exposures in the United States. *Exposure and Health*  
802 **15**, 373–394. <https://doi.org/10.1007/s12403-022-00496-y>
- 803 OECD (2022). Global Plastics Outlook: Policy Scenarios to 2060. Paris: OECD Publishing.  
804 <https://doi.org/10.1787/aa1edf33-en>
- 805 Pearce D, Atkinson G, Mourato S (2006) Cost-benefit analysis and the environment. Recent  
806 developments. Paris: OECD publications. <https://doi.org/10.1787/9789264010055-en>
- 807 Peng L, Fu D, Qi H, Lan CQ, Yu H and Ge C (2020). Micro-and nano-plastics in marine environment:  
808 Source, distribution and threats—A review. *Science of the total environment* **698**, 134254.  
809 <https://doi.org/10.1016/j.scitotenv.2019.134254>
- 810 Pew Charitable Trusts and Systemiq (2020). Breaking the plastic wave: A comprehensive assessment  
811 of pathways towards stopping plastic pollution. Available at:  
812 <https://www.systemiq.earth/breakingtheplasticwave/>
- 813 PlasticsEurope (2011) Plastics - the Facts 2011 An analysis of European plastics production, demand  
814 and recovery for 2010. Brussels: PlasticsEurope.
- 815 PlasticsEurope (2020) Plastics – the Facts 2020. An analysis of European plastics production, demand  
816 and waste data. Brussels: PlasticsEurope.
- 817 Ragusa A, Svelato A, Santacroce C, Catalano P, Notarstefano V, Carnevali O, ... and Giorgini E (2021)  
818 Plasticenta: First evidence of microplastics in human placenta. *Environment international* **146**, 106274.  
819 <https://doi.org/10.1016/j.envint.2020.106274>
- 820 Savoca MS, McInturf AG and Hazen EL (2021) Plastic ingestion by marine fish is widespread and  
821 increasing. *Global Change Biology* **27**(10), 2188-2199. <https://doi.org/10.1111/gcb.15533>
- 822 Science museum (2019) The age of plastic: from Parkesine to pollution. Web article published the 11<sup>th</sup>  
823 of October 2019. Available at: [https://www.sciencemuseum.org.uk/objects-and-stories/chemistry/age-](https://www.sciencemuseum.org.uk/objects-and-stories/chemistry/age-plastic-parkesine-pollution)  
824 [plastic-parkesine-pollution](https://www.sciencemuseum.org.uk/objects-and-stories/chemistry/age-plastic-parkesine-pollution)
- 825 Supran G, Rahmstorf S and Oreskes N (2023) Assessing ExxonMobil's global warming projections.  
826 *Science* **379**(6628), eabk0063. <https://doi.org/10.1126/science.abk0063>
- 827 The Ocean Cleanup (2021) Technology update. Conference presented at IMarEST annual conference  
828 (Institute of Marine Engineering, Science & Technology), July 5, Delft, The Netherlands. Available at:  
829 [https://www.youtube.com/watch?v=Wj\\_10gmQhPw](https://www.youtube.com/watch?v=Wj_10gmQhPw)

- 830 Timmer MP, Dietzenbacher E, Los B, Stehrer R, de Vries GJ (2015) An Illustrated User Guide to the  
831 World Input–Output Database: the Case of Global Automotive Production. *Review of International*  
832 *Economics* **23**, 575–605. <https://doi.org/10.1111/roie.12178>
- 833 Tjallema A (2022) Monitoring and performance evaluation of plastic cleanup systems developed by  
834 The Ocean Cleanup foundation. Conference presented at the 7<sup>th</sup> international marine debris conference  
835 (7IMDC), Busan, South Korea, 18-23 Septembre.
- 836 Trasande L, Zoeller RT, Hass U, Kortenkamp A, Grandjean P, Myers JP, ... and Heindel JJ (2015)  
837 Estimating burden and disease costs of exposure to endocrine-disrupting chemicals in the European  
838 Union. *The Journal of Clinical Endocrinology & Metabolism*, 100(4), 1245-1255.  
839 <https://doi.org/10.1210/jc.2014-4324>
- 840 Trasande L, Zoeller RT, Hass U, Kortenkamp A, Grandjean P, Myers JP, ... and Heindel JJ (2016)  
841 Burden of disease and costs of exposure to endocrine disrupting chemicals in the European Union: an  
842 updated analysis. *Andrology* **4**(4), 565-572. <https://doi.org/10.1111/andr.12178>
- 843 Trasande L (2022) A global plastics Treaty to protect endocrine health. *The Lancet Diabetes &*  
844 *Endocrinology* **10**(9), 616-618. [https://doi.org/10.1016/S2213-8587\(22\)00216-9](https://doi.org/10.1016/S2213-8587(22)00216-9)
- 845 Trasande L, Liu B, Bao W (2022a) Phthalates and attributable mortality: A population-based longitudinal  
846 cohort study and cost analysis. *Environmental Pollution* **292**(Part A), 118021.  
847 <https://doi.org/10.1016/j.envpol.2021.118021>
- 848 Uehara T, Cordier M and Hamaide B (2018) Fully dynamic input-output/system dynamics modeling for  
849 ecological-economic system analysis. *Sustainability* **10**(6), 1765. <https://doi.org/10.3390/su10061765>
- 850 UNEP (United Nations Environment Programme) (2014) Valuing Plastics: The Business Case for  
851 Measuring, Managing and Disclosing Plastic Use in the Consumer Goods Industry. Nairobi: UNEP  
852 publications. Available at: <https://wedocs.unep.org/20.500.11822/9238>
- 853 UNEP (United Nations Environment Programme) (2023) Turning off the Tap: How the world can end  
854 plastic pollution and create a circular economy. Nairobi: UNEP Publications. Available at:  
855 <https://www.unep.org/resources/turning-off-tap-end-plastic-pollution-create-circular-economy>
- 856 US EPA (2014) Guidelines for preparing economic analyses. 302 pp. Available at:  
857 [https://www.epa.gov/environmental-economics/guidelines-preparing-economic-analyses-](https://www.epa.gov/environmental-economics/guidelines-preparing-economic-analyses-2016#download)  
858 [2016#download](https://www.epa.gov/environmental-economics/guidelines-preparing-economic-analyses-2016#download)
- 859 US EPA (2020) Guidelines for preparing economic analyses. Review Copy Prepared for EPA's Science  
860 Advisory Board's Economic Guidelines Review Panel. 343 pp. Available at: [https://legacy-](https://legacy-assets.eenews.net/open_files/assets/2020/04/20/document_gw_02.pdf)  
861 [assets.eenews.net/open\\_files/assets/2020/04/20/document\\_gw\\_02.pdf](https://legacy-assets.eenews.net/open_files/assets/2020/04/20/document_gw_02.pdf)
- 862 van Emmerik T, Mellink Y, Hauk R, Waldschläger K and Schreyers L (2022) Rivers as plastic reservoirs.  
863 *Frontiers in Water* **3**, 786936. <https://doi.org/10.3389/frwa.2021.786936>
- 864 Vidal A (2020) Relocating rubbish. When Southeast Asia is overflowing with Western waste.  
865 *Visionscarto*. Available at: <https://visionscarto.net/relocating-rubbish>
- 866 Ward CP and Reddy CM (2020) We need better data about the environmental persistence of plastic  
867 goods. *Proceedings of the National Academy of Sciences* **117**(26), 14618-14621.  
868 <https://doi.org/10.1073/pnas.2008009117>
- 869 Wood B, Baker P, Scrinis G, McCoy D, Williams O and Sacks G (2021) Maximising the wealth of few  
870 at the expense of the health of many: a public health analysis of market power and corporate wealth and  
871 income distribution in the global soft drink market. *Globalization and Health* **17**, 1-17.  
872 <https://doi.org/10.1186/s12992-021-00781-6>

873 Yan H, Cordier M and Uehara T (2022) Demographic Factors and the Environmental Kuznets Curve:  
874 Global Plastic Pollution by 2050 Could Be 2 to 4 Times Worse than Projected. Preprint *SSRN 4231443*.  
875 Available at: [https://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=4231443](https://papers.ssrn.com/sol3/papers.cfm?abstract_id=4231443)

876



877 **Table 1. Global cost forecast of plastic pollution impacts (in case of inaction) and plastic pollution reduction**  
878 **strategies (in case of action towards zero plastics in ecosystems by 2040).** Note: all costs are in billion US\$ at  
879 prices of the year 2021 and are total values calculated over 2016-2040 with a discount rate of 3.5%. This Table is  
880 based on data from Sections 2 and 3.

	<b>COST TYPES</b>	<b>COST OF PLASTIC POLLUTION REDUCTION STRATEGIES</b>	<b>LOW ESTIMATE (US\$ BILLION)</b>	<b>HIGH ESTIMATE (US\$ BILLION)</b>
<b>Action scenario</b>	Private costs	Waste management costs *	470	1335
		<i>Terrestrial cleanup**</i>	507	1739
		Ocean cleanup (plastics floating offshore)**	11	248
		Ocean cleanup (plastics floating close to the shoreline)**	251	3895
		River cleanup (floating plastics)**	23	1373
		Cleanup of seabed, lakebed and riverbed (sinking plastics)	Omitted (due to lack of studies)	
		Cleanup of micro- and nano-plastics	Omitted (due to lack of studies)	
		Transition cost for the private sector towards 47% reduction of plastic production*	4847	5317
	External costs	Damages to marine ecosystems**	1003	132819
		Damages to terrestrial ecosystems	Omitted (due to lack of studies)	
		Human health in USA, EU and Canada **	11206	11692
		Human health in the rest of the world	Omitted (due to lack of studies)	
	<b>Social cost</b>	<b>Total cost of action</b>	<b>18318</b>	<b>158418</b>
<b>Inaction scenario</b>		<b>COST OF PLASTIC POLLUTION IMPACT</b>	<b>LOW ESTIMATE (US\$ BILLION)</b>	<b>HIGH ESTIMATE (US\$ BILLION)</b>
	Private cost	Waste management costs *	643	1612
	External costs	Damages to marine ecosystems**	1862	268498
		Damages to terrestrial ecosystems	Omitted (due to lack of studies)	
		Human health in USA, EU and Canada **	11206	11692
		Human health in the rest of the world	Omitted (due to lack of studies)	
	<b>Social cost</b>	<b>Total cost of inaction</b>	<b>13711</b>	<b>281802</b>
	<b>Comparison action/inaction</b>		Inaction (US\$ 13711 billion) is slightly cheaper than action (US\$ 18318 billion). However, given the costs and benefits calculated and the missing data (discussed in Section 5), it is not clear that the total cost of action is substantially higher than the one of inaction. Given the incomplete nature of this analysis, it is possible that the total cost of inaction is substantially higher as suggested by the high estimate in the last column of this table.	Inaction (US\$ 281802 billion) is significantly more expensive than action (US\$ 158418 billion)

881 \* Calculated in Section 2.2 for the “*system change scenario*”, which includes: (i) upstream interventions (reducing plastic  
882 production by 47% and substituting plastics with alternative materials), (ii) middle stream interventions (improving plastic waste  
883 collection and disposal, increasing plastic recycling), and a downstream solution (beach cleanup).

884 \*\* Calculated in Section 2.3 for cleanup of the legacy pollution, that is, plastic debris still remaining in terrestrial and aquatic  
885 ecosystems after implementing the “*system change scenario*”.

886 + Calculated in Section 2.2 for the BAU scenario.

887 ++ Calculated in Section 3.

888

889

890 **Table 2. Global benefits earned from plastic production in case of “Inaction” and “Action” scenarios**  
 891 **(scenarios described in Table 1).** Note: all benefits are in billion US\$ at prices of the year 2021 and are total  
 892 values calculated over 2016-2040 with a discount rate of 3.5%. Negative values are a cost. This table is based on  
 893 data from Sections 2, 3 and 4.

	<b>Benefits</b> (Obtained from plastic incomes: (taxes, wages & salaries, dividends, rents, etc.)		<b>Net benefit</b> (Benefits minus social costs calculated in table 1)	
	<b>Low estimate</b>	<b>High estimate</b>	<b>Low estimate</b>	<b>High estimate</b>
<b>Action scenario</b>	32668	33138	-120433	19667
<b>Inaction scenario</b>	37985	37985	-243817	24274
<b>Comparison action/inaction</b>	The “Action scenario” reduces incomes generated by plastic industries by 14% compared to the “Inaction scenario”	The “Action scenario” reduces incomes generated by plastic industries by 13 % compared to the “Inaction scenario”	<p>The net benefits in the “Action” and “Inaction” scenarios are both negative, which means an economic loss (that is, a cost).</p> <p>For the “Inaction scenario”, this means that the benefits obtained from the plastic industry are not sufficient to offset costs of plastic pollution impacts caused by inaction.</p> <p>For the “Action scenario”, the economic loss (that is, the negative net benefit) is significantly lower than in the “Inaction scenario”. This is because every year over 2021-2040, actions are implemented to reduce plastic pollution to approach the zero level in the ecosystems by 2040, which gradually reduces costs of plastic pollution impacts. These calculations should be repeated in further studies, when more data on costs and benefits become available, in order to check whether the low estimate of the net benefit of the “Action scenario” becomes positive.</p>	<p>Net benefits earned in the “Action” and “Inaction” scenarios are both positive, which represents an economic gain.</p> <p>For the “Action scenario”, this suggests that actions towards zero plastics pollution by 2040 is profitable for society because reduced cost of damages resulting from plastic pollution reduction strategies are sufficient to offset costs of actions.</p> <p>The net benefit in the “Inaction scenario” is slightly higher than in the “Action scenario”. This is because in the calculations of the “Inaction scenario”, production is not reduced and, hence, benefits obtained from the plastic industry seem to more than compensate costs of plastic pollution impacts caused by inaction. However, given the incomplete nature of this analysis (several cost and benefit data are lacking as discussed in Section 5), it is not clear that the net benefit of inaction is substantially higher than the one of action. On the contrary, when more data will be made available, further studies might show it is possible that the net benefit of inaction is substantially lower than the one of action.</p>


894

895

## 896 List of figures

897 – Figure 1. Global cumulative discard of plastic waste inadequately managed over 1950-2060 – BAU  
 898 scenario. Note: MMT: million metric tons. The curves are computed summing over time global annual discard of  
 899 inadequately managed plastic waste (Figure S1, in Supplemental materials) provided by Lebreton and Andrady (2019),  
 900 Cordier et al. (2021), Yan et al. (2022) and Lau et al. (2020).


**Global accumulated discard of inadequately managed plastic waste (MMT)**



901

902

903 – Figure 2. Global plastic debris accumulated over time in terrestrial (upper graph) and aquatic (lower  
 904 graph) ecosystems over 1950-2060 – BAU scenario. Note: aquatic ecosystems include lakes, rivers and oceans  
 905 globally. The curves are obtained summing over time annual emissions of plastic waste into the ecosystems (Figure S2,  
 906 in Supplemental materials) provided by Borrelle et al. (2020), Lau et al. (2020) and the OECD (2022). The OECD (2022)  
 907 also provides accumulated values in 2019 and 2060. We used them to cross-check our computation method and make sure  
 908 we did not make any mistakes.




909

910




911 – Figure 3. Global plastic debris accumulated over time in aquatic ecosystems disaggregated into oceans  
 912 (upper graph), plastics floating in rivers (middle graph), and plastics sinking on riverbeds and lakebeds  
 913 (lower graph) – BAU scenario. Note: the curves are obtained summing over time estimations of annual emissions  
 914 of plastic waste (Figure S3, in Supplemental materials) provided by Lebreton et al. (2019). The other models directly  
 915 provided accumulated values (Jambeck et al., 2015; Cordier and Uehara, 2019; and OECD, 2022).



916

917


918 – Figure 4. Comparison of global total cost of action (left bars) and inaction (right bars) over 2016-  
919 2040. Note: the graph is based on data from Table 1. The lower estimates suggest the cost of inaction (US\$ 13711 billion)  
920 is slightly cheaper than the one of action (US\$ 18318 billion). However, given the costs and benefits calculated and the  
921 missing data (discussed in Section 5), it is not clear that the total cost of action is substantially higher than the one of  
922 inaction. Given the incomplete nature of this analysis, it is possible that the total cost of inaction is substantially higher as  
923 suggested by the high estimate (inaction cost: US\$ 281802 billion, which is significantly more expensive than action cost:  
924 US\$ 158418 billion).




925

926

Figure 5. Comparison of global total net benefit of action (left bars) and inaction (right bars) over 2016-2040. Note: the graph is based on data from Table 2. Net benefit = benefit earned from plastics minus costs. The low estimate of net benefits in the “Action” and “Inaction” scenarios are both negative, which means an economic loss (that is, a cost). For the “Inaction scenario”, this means that the benefits obtained from the plastic industry are not sufficient to offset costs of plastic pollution impacts caused by inaction. For the “Action scenario”, the economic loss (that is, the negative net benefit) is significantly lower than in the “Inaction scenario”. This is because every year over 2021-2040, actions are implemented to reduce plastic pollution to approach the zero level in the ecosystems by 2040, which gradually reduces costs of plastic pollution impacts. These calculations should be repeated in further studies, when more data on costs and benefits become available (see missing data listed in Table 1), in order to check whether the low estimate of the net benefit of the “Action scenario” becomes positive. The high estimate of net benefits earned in the “Action” and “Inaction” scenarios are both positive, which represents an economic gain. For the “Action scenario”, this suggests that actions towards zero plastics pollution by 2040 is profitable for society because reduced cost of damages resulting from plastic pollution reduction strategies are sufficient to offset costs of actions. The high estimate of the net benefit in the “Inaction scenario” is slightly higher than in the “Action scenario”. This is because in the calculations of the “Inaction scenario”, production is not reduced and, as a result, the benefits obtained from the plastics industry appear to more than offset the costs of the impacts of plastic pollution caused by inaction. However, given the incomplete nature of this analysis (several cost and benefit data are lacking as discussed in Section 5), it is not clear that the high estimate of the net benefit of inaction is substantially higher than the one of action. On the contrary, when more data will be made available, further studies might show it is possible that the net benefit of inaction is substantially lower than the one of action.



948 – Figure 6. Cost of plastic pollution reduction policies as simulated in the global ambitious policy  
 949 scenario by the OECD (2022, p. 198). Note: costs are expressed as a percentage of GDP (Gross Domestic Product).  
 950 World regions that are part of the Global South are in red and Global North regions are in blue. The black bar shows the  
 951 world average cost (average calculated across all countries).



952

953



Photograph by Juan Baztan, CEARC-UVSQ, Zero Plastic in Biosphere Reserves, cc-by-sa-nc.