



HAL
open science

Denominator vectors and dimension vectors from triangulated surfaces

Toshiya Yurikusa

► **To cite this version:**

Toshiya Yurikusa. Denominator vectors and dimension vectors from triangulated surfaces. *Journal of Algebra*, 2024, 641, pp.620-647. 10.1016/j.jalgebra.2023.12.002 . hal-04439601

HAL Id: hal-04439601

<https://hal.uvsq.fr/hal-04439601>

Submitted on 30 May 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DENOMINATOR VECTORS AND DIMENSION VECTORS FROM TRIANGULATED SURFACES

TOSHIYA YURIKUSA

ABSTRACT. In a categorification of skew-symmetric cluster algebras, each cluster variable corresponds with an indecomposable module over the associated Jacobian algebra. Buan, Marsh and Reiten studied when the denominator vector of each cluster variable in an acyclic cluster algebra coincides with the dimension vector of the corresponding module. In this paper, we give analogues of their results for cluster algebras from triangulated surfaces by comparing two kinds of intersection numbers of tagged arcs.

1. INTRODUCTION

Cluster algebras [FZ02] are commutative algebras with generators called cluster variables. The certain tuples of cluster variables are called clusters. In a cluster algebra with principal coefficients, by Laurent phenomenon, any cluster variable x is expressed by a Laurent polynomial of the initial cluster variables (x_1, \dots, x_n) and coefficients (y_1, \dots, y_n)

$$x = \frac{F(x_1, \dots, x_n, y_1, \dots, y_n)}{x_1^{d_1(x)} \dots x_n^{d_n(x)}},$$

where $d_i(x) \in \mathbb{Z}$ and $F(x_1, \dots, x_n, y_1, \dots, y_n) \in \mathbb{Z}[x_1, \dots, x_n, y_1, \dots, y_n]$ is not divisible by any x_i [FZ02, FZ07]. We call $d(x) := (d_i(x))_{1 \leq i \leq n}$ the denominator vector of x . For the maximal degree $f_i(x)$ of y_i in the F -polynomial $F(1, \dots, 1, y_1, \dots, y_n)$, we call $f(x) := (f_i(x))_{1 \leq i \leq n}$ the f -vector of x . It was conjectured that $d(x) = f(x)$ for any non-initial cluster variable x in [FZ07, Conjecture 7.17].

Let Q be a quiver without oriented cycles of length at most two and K an algebraically closed field. The quiver Q defines a cluster algebra $\mathcal{A}(Q)$ [FZ02] and a Jacobian algebra $J(Q) = J(Q, W)$ over K for a non-degenerate potential W of Q [DWZ08]. Then non-initial cluster variables x of $\mathcal{A}(Q)$ correspond with certain indecomposable $J(Q)$ -modules $M(x)$ (see Subsection 3.2). Note that the dimension vector $\underline{\dim} M(x)$ is independent of the choice of W as stated below. It was proved in [CK06] (see also [CC06, Hub06]) that if Q is acyclic, then it satisfies the property

$$(d=\dim) \quad d(x) = \underline{\dim} M(x) \text{ for any non-initial cluster variable } x \text{ of } \mathcal{A}(Q).$$

However, it does not hold in general though it is known that $d(x) \leq \underline{\dim} M(x)$ [DWZ10, Corollary 5.5]. Buan, Marsh, and Reiten [BMR09] gave a characterization of quivers satisfying $(d=\dim)$ for mutation equivalence classes of acyclic quivers (see also [GLFS16, Theorem 1.1]).

Theorem 1.1 ([BMR09, Theorems 1.5 and 1.6]). *Let Q be a quiver obtained from an acyclic quiver by a sequence of mutations. If Q satisfies $(d=\dim)$, then $\text{End}_{J(Q)}(P) \simeq K$ for any indecomposable projective $J(Q)$ -module P . In addition, if $J(Q)$ is a tame algebra, then the inverse holds.*

Note that [BMR09, Theorems 1.5 and 1.6] were given in terms of the associated cluster category, but they are equivalent to Theorem 1.1 by [BMR09, Remark 3.4] (see (3.1)). They also gave the following result as a consequence of Theorem 1.1.

Theorem 1.2 ([BMR09, Corollary 1.7]). *Let Q be an acyclic quiver. Then every quiver obtained from Q by a sequence of mutations satisfies $(d=\dim)$ if and only if Q is a Dynkin quiver or has exactly two vertices.*

2020 Mathematics Subject Classification. 13F60, 16G20.

Key words and phrases. Cluster algebra, marked surface, denominator, dimension vector.

On the other hand, in the case that $J(Q)$ is finite dimensional, Fu and Keller [FK10] proved that the equality $f(x) = \underline{\dim} M(x)$ holds and it gave a counterexample of [FZ07, Conjecture 7.17] (based on [BMR09]). This equality holds in general (Theorem 3.8), thus the vector $\underline{\dim} M(x)$ and the property ($d=\dim$) are independent of the choice of W since f -vectors are so.

In this paper, we consider Theorems 1.1 and 1.2 for quivers defined from triangulated surfaces. Cluster algebras associated with marked surfaces were introduced in [FST08] based on [FG06]. To a tagged triangulation T of a marked surface \mathcal{S} , they associated a quiver Q_T and studied the cluster algebra $\mathcal{A}(Q_T)$. In particular, tagged arcs (resp., tagged triangulations, tagged arcs of T) of \mathcal{S} correspond bijectively with cluster variables (resp., clusters, the initial cluster variables) of $\mathcal{A}(Q_T)$ except that \mathcal{S} is a closed surface with exactly one puncture. Moreover, flips of tagged triangulations are compatible with mutations of clusters (see Section 4).

Moreover, there are two kinds of intersection numbers of tagged arcs: One was introduced in [FST08] to give denominator vectors in $\mathcal{A}(Q_T)$; Another one was introduced in [QZ17] to give dimension of homomorphisms in the associated cluster category. Then the latter also gives f -vectors in $\mathcal{A}(Q_T)$ (Theorem 4.2). Therefore, we mainly study these intersection numbers. As a consequence, we characterize quivers satisfying ($d=\dim$) in geometric terms.

Theorem 1.3. *Let \mathcal{S} be a marked surface and T a tagged triangulation of \mathcal{S} . Then Q_T satisfies ($d=\dim$) if and only if either \mathcal{S} is a torus with exactly one puncture or T has neither loops nor tagged arcs connecting punctures.*

Theorem 1.4. *Let \mathcal{S} be a marked surface.*

- (1) *There is at least one tagged triangulation T of \mathcal{S} such that Q_T satisfies ($d=\dim$) if and only if \mathcal{S} is one of the following:*
 - *a marked surface of genus zero with non-empty boundary;*
 - *a marked surface of positive genus with non-empty boundary and at least three marked points, at least two of which are on $\partial\mathcal{S}$;*
 - *a torus with exactly one puncture.*
- (2) *The quivers Q_T satisfy ($d=\dim$) for all tagged triangulations T of \mathcal{S} if and only if \mathcal{S} is one of the following:*
 - *a polygon with at most one puncture;*
 - *an annulus with exactly two marked points;*
 - *a torus with exactly one puncture.*

An analogue of Theorem 1.2 immediately follows from Theorem 1.4(2) (see Example 4.1), where we say that a quiver is Dynkin type if it is obtained from a Dynkin quiver by a sequence of mutations.

Corollary 1.5. *Let T be a tagged triangulation. Then every quiver obtained from Q_T by a sequence of mutations satisfies ($d=\dim$) if and only if Q_T is either Dynkin type, a Kronecker quiver, or a Markov quiver.*

Moreover, we also give an analogue of Theorem 1.1 except for closed surfaces with exactly one puncture by using Theorem 1.3.

Theorem 1.6 (Theorem 4.7). *Let T be a tagged triangulation of a marked surface which is not a closed surface with exactly one puncture. Then Q_T satisfies ($d=\dim$) if and only if $\text{End}_{J(Q_T)}(P) \simeq K$ for any indecomposable projective $J(Q_T)$ -module P .*

Finally, if \mathcal{S} is a torus with exactly one puncture, then Q_T satisfies ($d=\dim$) by Theorem 1.3. However, Labardini-Fragoso [LF09] constructed a non-degenerate potential W of Q_T such that

- $\dim \text{End}_{J(Q_T, W)}(P) = 4$ for any indecomposable projective $J(Q_T, W)$ -module P [Lad12, Proposition 4.2];
- $J(Q_T, W)$ is tame [GLFS16] (see also [LF16a, Theorem 3.5]).

Therefore, Theorem 1.1 (or Theorem 1.6) does not hold in general.

This paper is organized as follows. In Section 2, we only work on marked surfaces. We study two kinds of intersection numbers and give geometric versions of Theorems 1.3 and 1.4 (Theorems 2.10, 2.11 and 2.12). In Section 3, we recall cluster algebras and their categorification. For the case arising from marked surfaces, they have many properties as in Section 4. Using the properties and the results in Section 2, we give Theorems 1.3, 1.4 in Subsection 4.1 and Theorem 1.6 in Subsection 4.2. Moreover, we also characterize the condition of Corollary 1.5 in terms of the associated cluster category (Proposition 4.9). In Section 5, we give an example which is not covered by [BMR09].

2. ON MARKED SURFACES

In this section, we work on marked surfaces [FST08, FT18].

2.1. Intersection numbers of tagged arcs. Let S be a connected compact oriented Riemann surface with (possibly empty) boundary ∂S and M a non-empty finite set of marked points on S with at least one marked point on each connected component of ∂S . We call the pair $\mathcal{S} = (S, M)$ a *marked surface*. A marked point in the interior of S is called a *puncture*. For technical reasons, we assume that \mathcal{S} is not a monogon with at most one puncture, a digon without punctures, a triangle without punctures, and a sphere with at most three punctures (see [FST08] for the details). Throughout this paper, we consider a curve on S up to isotopy relative to M .

A *tagged arc* of \mathcal{S} is a curve on S whose endpoints are in M and each end is tagged in one of two ways, *plain* or *notched*, such that the following conditions are satisfied:

- it does not intersect itself except at its endpoints;
- it is disjoint from M and ∂S except at its endpoints;
- it does not cut out a monogon with at most one puncture or a digon without punctures;
- its ends incident to ∂S are tagged plain;
- both ends of a loop are tagged in the same way,

where a *loop* is a tagged arc with two identical endpoints. In the figures, we represent tags as follows:

plain —● notched —⊗● .

A *pair of conjugate arcs* is the following pair of tagged arcs whose underlying curves coincide:



We denote by $\mathbb{A}_{\mathcal{S}}$ the set of tagged arcs of \mathcal{S} . We also recall a rotation of tagged arcs [BQ15] (see Table 2).

Definition 2.1. The *tagged rotation* of $\gamma \in \mathbb{A}_{\mathcal{S}}$ is the tagged arc $\rho(\gamma)$ defined as follows:

- If γ has an endpoint o on a component C of ∂S , then $\rho(\gamma)$ is obtained from γ by moving o to the next marked point on C in the counterclockwise direction;
- If γ has an endpoint at a puncture p , then $\rho(\gamma)$ is obtained from γ by changing its tags at p .

Throughout this paper, when we consider intersections of curves, we assume that they intersect transversally in a minimum number of points in $S \setminus M$. Our main subjects are two kinds of intersection numbers of tagged arcs defined in [FST08, Definition 8.4] and [QZ17, Definition 3.3].

Definition 2.2. Let $\gamma, \delta \in \mathbb{A}_{\mathcal{S}}$.

- (1) The *FST-intersection number* $(\gamma|\delta)$ of γ and δ is defined by $A_{\gamma,\delta} + B_{\gamma,\delta} + C_{\gamma,\delta} + D_{\gamma,\delta}$, where
 - $A_{\gamma,\delta}$ is the number of intersection points of γ and δ in $S \setminus M$;
 - $B_{\gamma,\delta} = 0$ unless γ is a loop at $o \in M$, in which case $B_{\gamma,\delta}$ is the negative of the number of contractible triangles Δ defined as follows: Suppose that δ consecutively intersects with γ at points a and b . Then Δ consists of the segment of δ from a to b , the segment of γ from o to a , and the segment of γ from o to b (see Figure 1);
 - $C_{\gamma,\delta} = 0$ unless the underlying curves of γ and δ coincide, in which case $C_{\gamma,\delta} = -1$;

- $D_{\gamma,\delta}$ is the number of ends of δ incident to an endpoint e of γ whose tags are different from one of γ at e .
- (2) The *QZ-intersection number* $\text{Int}(\gamma, \delta)$ of γ and δ is defined by $A_{\gamma,\delta} + C'_{\gamma,\delta} + D'_{\gamma,\delta}$, where
- $C'_{\gamma,\delta} = 0$ unless γ and δ form a pair of conjugate arcs, in which case $C'_{\gamma,\delta} = -1$;
 - $D'_{\gamma,\delta}$ is the number of pairs of an end of γ and an end of δ such that they are incident to a common puncture and their tags are different.




FIGURE 1. The triangle Δ contributes -1 to B if it is contractible

Note that FST-intersection numbers are not symmetric in general, while QZ-intersection numbers are symmetric.

Let $\gamma, \delta \in \mathbb{A}_{\mathcal{S}}$. By the definitions, $C_{\gamma,\delta}$, $D_{\gamma,\delta}$, $C'_{\gamma,\delta}$, and $D'_{\gamma,\delta}$ are zero if γ and δ have no common endpoints. Moreover, $C_{\gamma,\delta}$ and $C'_{\gamma,\delta}$ are zero if the underlying curves of γ and δ do not coincide. Thus Table 1 lists all cases where any of them are not zero up to changing all tags at each puncture. In particular, $(\gamma|\gamma) = C_{\gamma,\gamma} = -1 \neq 0 = C'_{\gamma,\gamma} = \text{Int}(\gamma, \gamma)$. Moreover, if γ and δ form a pair of conjugate arcs, then $(\gamma|\delta) = \text{Int}(\gamma, \delta) = 0$.

$C_{\gamma,\delta}$	-1	-1	-1			
$C'_{\gamma,\delta}$	0	-1	0			
$C''_{\gamma,\delta}$	-1	0	-1			
$D_{\gamma,\delta}$	1	2	1	1	2	2
$D'_{\gamma,\delta}$	1	2	1	2	2	4
$D''_{\gamma,\delta}$	0	0	0	-1	0	-2

TABLE 1. Cases that any of $C_{\gamma,\delta}$, $D_{\gamma,\delta}$, $C'_{\gamma,\delta}$, and $D'_{\gamma,\delta}$ are not zero

By the definitions, we can give a difference between the intersection numbers.

Proposition 2.3. *Let $\gamma, \delta \in \mathbb{A}_{\mathcal{S}}$. Then we have*

$$(\gamma|\delta) = \text{Int}(\gamma, \delta) + B_{\gamma,\delta} + C''_{\gamma,\delta} + D''_{\gamma,\delta},$$

where

- $B_{\gamma,\delta}$ is defined in Definition 2.2(1);
- $C''_{\gamma,\delta} = 0$ unless the underlying curves of γ and δ coincide and they do not form a pair of conjugate arcs, in which case $C''_{\gamma,\delta} = -1$;
- $D''_{\gamma,\delta} = 0$ unless γ is a loop at a puncture p , in which case $D''_{\gamma,\delta}$ is the negative of the number of ends of δ incident to p whose tags are different from ones of γ .

Proof. Table 1 gives the equalities $C''_{\gamma,\delta} = C_{\gamma,\delta} - C'_{\gamma,\delta}$ and $D''_{\gamma,\delta} = D_{\gamma,\delta} - D'_{\gamma,\delta}$. Otherwise, all of them are zero as the above observation. Therefore, we get the desired equality from Definition 2.2. \square

We observe some properties of the values $B_{\gamma,\delta}$, $C''_{\gamma,\delta}$, and $D''_{\gamma,\delta}$.

Lemma 2.4. *Let $\gamma, \delta \in \mathbb{A}_{\mathcal{S}}$.*

- (1) *The values $B_{\gamma,\delta}$, $C''_{\gamma,\delta}$, and $D''_{\gamma,\delta}$ are non-positive and $(\gamma|\delta) \leq \text{Int}(\gamma, \delta)$.*
- (2) *If γ is not a loop, then $B_{\gamma,\delta} = D''_{\gamma,\delta} = 0$.*
- (3) *There is a $\gamma' \in \mathbb{A}_{\mathcal{S}} \setminus \{\gamma\}$ such that $C''_{\gamma,\gamma'} \neq 0$ if and only if γ connects punctures.*

Proof. The assertions immediate follow from the definitions. \square

Lemma 2.5. *If γ is a loop of \mathcal{S} at $m \in M \cap \partial\mathcal{S}$, then there is a $\delta \in \mathbb{A}_{\mathcal{S}}$ such that $B_{\gamma,\delta} \neq 0$.*

Proof. Let C be a connected component of $\partial\mathcal{S}$ containing m . If γ is homotopic to C , then there are other marked points on C . We take a loop δ at such a marked point which is homotopic to C . Then we have $B_{\gamma,\delta} \neq 0$ (see the left diagram of Figure 2).

Suppose that γ is not homotopic to C . We take δ as a tagged arc obtained from γ by applying the Dehn twist along a simple closed curve which is homotopic to C (see e.g. [Mar16] for the definition of Dehn twists). Then we get $B_{\gamma,\delta} \neq 0$ (see the right diagrams of Figure 2). \square





FIGURE 2. Constructions of $\delta \in \mathbb{A}_{\mathcal{S}}$ satisfying $B_{\gamma,\delta} \neq 0$ for a loop γ at $m \in M \cap \partial\mathcal{S}$

Lemma 2.6. *Suppose that \mathcal{S} is a closed surface with exactly one puncture. Let $\gamma \in \mathbb{A}_{\mathcal{S}}$.*


- (1) *If the genus of \mathcal{S} is one, then we have $B_{\gamma,\delta} = 0$ for any $\delta \in \mathbb{A}_{\mathcal{S}}$.*
- (2) *If the genus of \mathcal{S} is two or more, then there is a $\delta \in \mathbb{A}_{\mathcal{S}}$ such that $B_{\gamma,\delta} \neq 0$.*

Proof. The claim (1) was proved in [RS17, Subsection 4.3] in the case that all ends of γ and δ are tagged plain. Since $B_{\gamma,\delta}$ is independent of their tags, it holds in general cases.


For (2), we can take a desired δ concretely. In fact, we consider the case that the genus of \mathcal{S} is two. Then \mathcal{S} is described by the octahedron P_8 as follows:



where c_1, \dots, c_8 are segments of a simple closed curve which is contracted to a puncture, and the edges of P_8 with the same numbers are identified along the directions. In P_8 , there are the connected segments γ' and γ'' of γ containing endpoints of γ , where they may be identified. We assume that the endpoints of γ' and γ'' are p_i and p_j for $i \leq j$, respectively. Then γ' and γ'' intersect with c_i and c_j , and let q_i and q_j be their intersection points, respectively. If δ contains the curve $c_i c_{i+1} \cdots c_j$ or $c_j \cdots c_n c_1 \cdots c_i$ in \mathcal{S} , then δ consecutively intersects with γ at q_i and q_j . Then the segment of δ from q_i to q_j , the segment of γ from p to q_i , and the segment of γ from p to q_j form a contractible triangle, thus we have $B_{\gamma, \delta} \neq 0$. That is, we only need to give such δ . We consider four tagged arcs



Then δ_k contains $c_{k+1} \cdots c_{k+6}$ for $k \in \{1, 2, 5, 6\}$, where $c_{8+h} = c_h$. Therefore, there is k such that $B_{\gamma, \delta_k} \neq 0$ except for $(i, j) = (1, 5)$. In which case, either $B_{\gamma, \delta_3} \neq 0$ or $B_{\gamma, \delta_7} \neq 0$ for two tagged arcs



In the same way, we can construct a desired δ for the case that the genus of S is three or more. \square

2.2. Tagged triangulations with common intersection vectors. A *tagged triangulation* of \mathcal{S} is a maximal set T of tagged arcs of \mathcal{S} such that $\text{Int}(\gamma, \delta) = 0$ for any $\gamma, \delta \in T$. We denote by $\mathbb{T}_{\mathcal{S}}$ the set of tagged triangulations of \mathcal{S} . By [FST08, Remark 4.2], a tagged triangulation of \mathcal{S} decomposes \mathcal{S} into *puzzle pieces* as in Figure 3.




FIGURE 3. A complete list of puzzle pieces of \mathcal{S} up to changing all tags at each puncture, where the rightmost puzzle piece forms a tagged triangulation of a sphere with exactly four punctures

Let $T \in \mathbb{T}_{\mathcal{S}}$. For $\gamma \in T$, there is a unique tagged arc $\gamma' \notin T$ such that $\mu_{\gamma} T := (T \setminus \{\gamma\}) \sqcup \{\gamma'\} \in \mathbb{T}_{\mathcal{S}}$. We call $\mu_{\gamma} T$ the *flip* of T at γ . We denote by $\mathbb{T}_{\mathcal{S}}^T$ the set of tagged triangulations of \mathcal{S} obtained from T by sequences of flips, and by $\mathbb{A}_{\mathcal{S}}^T$ the set of tagged arcs of tagged triangulations contained in $\mathbb{T}_{\mathcal{S}}^T$. It is elementary that flips are compatible with tagged rotations, that is, we have $\rho(\mu_{\gamma} T) = \mu_{\rho(\gamma)} \rho(T)$. It implies that $\rho : \mathbb{T}_{\mathcal{S}}^T \rightarrow \mathbb{T}_{\mathcal{S}}^{\rho(T)}$ is a bijection which commutes with flips. In almost cases, its domain and codomain coincide as follows.

Theorem 2.7 ([FST08, Theorem 7.9 and Proposition 7.10]). *If \mathcal{S} is not a closed surface with exactly one puncture, then we have $\mathbb{T}_{\mathcal{S}}^T = \mathbb{T}_{\mathcal{S}}$, that is, any two tagged triangulations of \mathcal{S} are connected by a*

sequence of flips. If \mathcal{S} is a closed surface with exactly one puncture, then we have $\mathbb{T}_{\mathcal{S}}^T \sqcup \mathbb{T}_{\mathcal{S}}^{\rho(T)} = \mathbb{T}_{\mathcal{S}}$, in particular, all tags appearing in $\mathbb{T}_{\mathcal{S}}^T$ are the same.

For $\gamma \in \mathbb{A}_{\mathcal{S}}$, we set two kinds of *intersection vectors*

$$(T|\gamma) := ((t|\gamma))_{t \in T} \in \mathbb{Z}^T \quad \text{and} \quad \text{Int}(T, \gamma) := (\text{Int}(t, \gamma))_{t \in T} \in \mathbb{Z}^T.$$

Note that the former induces the denominator vector of the corresponding cluster variable, while the latter induces its f -vector and the dimension vector of the corresponding indecomposable module (see Theorem 4.2 and under Theorem 3.8).

Definition 2.8. Let $T \in \mathbb{T}_{\mathcal{S}}$. We say that

- (1) T has *common intersection vectors* (CIV, for short) if $(T|\gamma) = \text{Int}(T, \gamma)$ for any $\gamma \in \mathbb{A}_{\mathcal{S}} \setminus T$;
- (2) T has *weak common intersection vectors* (wCIV, for short) if $(T|\gamma) = \text{Int}(T, \gamma)$ for any $\gamma \in \mathbb{A}_{\mathcal{S}}^T \setminus T$.

Theorem 2.7 implies that CIV and wCIV are equivalent if \mathcal{S} is not a closed surface with exactly one puncture. In which case, we get the following.

Proposition 2.9. *Suppose that \mathcal{S} is a closed surface with exactly one puncture. Then all $T \in \mathbb{T}_{\mathcal{S}}$ do not have CIV. Moreover, the following are equivalent:*

- (1) *There is at least one $T \in \mathbb{T}_{\mathcal{S}}$ with wCIV;*
- (2) *All $T \in \mathbb{T}_{\mathcal{S}}$ have wCIV;*
- (3) *\mathcal{S} is a torus.*

Proof. Using the notations in Proposition 2.3, the first claim follows from $C''_{\gamma, \rho(\gamma)} \neq 0$ for any $\gamma \in \mathbb{A}_{\mathcal{S}}$ under the assumption. Let $T \in \mathbb{T}_{\mathcal{S}}$ and $\gamma \in T$. We have $C''_{\gamma, \delta} = D''_{\gamma, \delta} = 0$ for any $\delta \in \mathbb{A}_{\mathcal{S}}^T \setminus \{\gamma\}$ since all tags of γ and δ are the same by Theorem 2.7. By Lemma 2.6, there is a $\delta \in \mathbb{A}_{\mathcal{S}}$ such that $B_{\gamma, \delta} \neq 0$ if and only if \mathcal{S} is not a torus. In particular, we can take δ as in $\mathbb{A}_{\mathcal{S}}^T$ since $B_{\gamma, \delta}$ is independent of tags. Therefore, T have wCIV if and only if \mathcal{S} is a torus. This gives the desired equivalences. \square

We are ready to give the main result of this section.

Theorem 2.10. *For $T \in \mathbb{T}_{\mathcal{S}}$, the following are equivalent:*

- (1) *T has CIV;*
- (2) *T has neither loops nor tagged arcs connecting punctures;*
- (3) *$\text{Int}(\gamma, \rho(\gamma)) = 1$ for any $\gamma \in T$;*
- (4) *$\text{Int}(\gamma, \rho^{-1}(\gamma)) = 1$ for any $\gamma \in T$.*

Proof. Using the notations in Proposition 2.3, Lemma 2.4(1) implies that T have CIV if and only if $B_{\gamma, \delta} = C''_{\gamma, \delta} = D''_{\gamma, \delta} = 0$ for any $\gamma \in T$ and $\delta \in \mathbb{A}_{\mathcal{S}} \setminus \{T\}$. By Lemma 2.4(2) and (3), their equalities hold for T satisfying (2), that is, (2) implies (1). The inverse follows from Lemmas 2.4(3) and 2.5. It is straightforward to check the remaining equivalences. In fact, Table 2 lists all kinds of tagged arcs up to changing all tags at each puncture. It gives that $\gamma \in \mathbb{A}_{\mathcal{S}}$ is neither a loop nor a tagged arc connecting punctures if and only if $\text{Int}(\gamma, \rho(\gamma)) = 1$. Similarly, it is equivalent to $\text{Int}(\gamma, \rho^{-1}(\gamma)) = 1$. Thus (2), (3) and (4) are equivalent. \square

Theorem 2.11. *For $T \in \mathbb{T}_{\mathcal{S}}$, it has wCIV if and only if it has CIV or \mathcal{S} is a torus with exactly one puncture.*

Proof. Since CIV and wCIV are equivalent except that \mathcal{S} is a closed surface with exactly one puncture, the assertion follows from Proposition 2.9. \square

In general, the conditions (1) and (2) in Proposition 2.9 are not equivalent. We also characterize \mathcal{S} satisfying each condition.

Theorem 2.12. *Let \mathcal{S} be a marked surface.*

- (1) *There is at least one $T \in \mathbb{T}_{\mathcal{S}}$ with CIV if and only if \mathcal{S} is one of the following:*

$\text{Int}(\gamma, \rho(\gamma))$	1	1	2	2	4

TABLE 2. QZ-intersection number of each tagged arc γ and its tagged rotation $\rho(\gamma)$

- a marked surface of genus zero with $\partial S \neq \emptyset$;
 - a marked surface of positive genus with $\partial S \neq \emptyset$, $|M| \geq 3$ and $|M \cap \partial S| \geq 2$.
- (2) All $T \in \mathbb{T}_{\mathcal{S}}$ have CIV if and only if \mathcal{S} is one of the following:
- a polygon with at most one puncture;
 - an annulus with $|M| = 2$.

In (1) and (2), we also have the similar statements obtained by replacing CIV with $w\text{CIV}$ and adding the case that \mathcal{S} is a torus with exactly one puncture.

Proof of Theorem 2.12(2). By Theorem 2.10, all $T \in \mathbb{T}_{\mathcal{S}}$ have CIV if and only if there are neither loops nor tagged arcs connecting punctures in \mathcal{S} . It is easy to check that \mathcal{S} appearing in (2) satisfies such conditions. In other cases, \mathcal{S} satisfies at least one of the following:

- there are at least two punctures;
- the genus of S is positive;
- there are at least two boundary components and $|M| \geq 3$.

In each case, we can take a loop or a tagged arc connecting punctures as in Figure 4. Thus (2) holds. \square




FIGURE 4. A loop or a tagged arc connecting punctures on each marked surface which does not appear in Theorem 2.12(2)


To prove Theorem 2.12(1), we prepare the following property.

Proposition 2.13. *Suppose that $\partial S \neq \emptyset$ and there is a $T \in \mathbb{T}_{\mathcal{S}}$ with CIV. Then there is a $T' \in \mathbb{T}_{\mathcal{S}'}$ with CIV if \mathcal{S}' is a marked surface obtained from \mathcal{S} by one of the following:*

- (1) adding a marked point on ∂S ;
- (2) adding a puncture;
- (3) adding a boundary component with exactly one marked point.

Proof. Up to homotopy, we can take a puzzle piece Δ of T whose side is a boundary segment of \mathcal{S} such that the added marked point or boundary component is in Δ . We obtain $T' \in \mathbb{T}_{\mathcal{S}'}$ from T by

adding puzzle pieces in each case as follows:




In all cases, T' has neither loops nor tagged arcs connecting punctures since T is so. Thus it also has CIV by Theorem 2.10. \square

Proof of Theorem 2.12(1). If $\partial S = \emptyset$, then any tagged arc of \mathcal{S} connects punctures. Thus any $T \in \mathbb{T}_{\mathcal{S}}$ does not have CIV by Theorem 2.10. Therefore, we assume that $\partial S \neq \emptyset$. Let g be the genus of S and P the set of punctures of \mathcal{S} . We consider five cases.

Case 1. $g = 0$. By Theorem 2.12(2) and Proposition 2.13, there is a $T \in \mathbb{T}_{\mathcal{S}}$ with CIV except for a monogon with at least two punctures since we assume that \mathcal{S} is not a monogon with at most one puncture. For a monogon with exactly two punctures, two pairs of conjugate arcs form a tagged triangulation with CIV as in the proof of Proposition 2.13. By Proposition 2.13 again, there is a $T \in \mathbb{T}_{\mathcal{S}}$ with CIV for any \mathcal{S} with $g = 0$.

Case 2. $g \geq 1, |M \setminus P| = 1$. Suppose that there is a $T \in \mathbb{T}_{\mathcal{S}}$ with CIV, that is, T has neither loops nor tagged arcs connecting punctures by Theorem 2.10. By this condition, T must be constructed as follows:




In fact, for the unique boundary segment b , a puzzle piece of T with b must be a triangle brl_1 whose one vertex is a puncture. If $l_1 = r$, then S is a sphere, a contradiction. Suppose that $l_1 \neq r$. Then another puzzle pieces of T with l_1 must be a digon l_1l_2 . Similarly, a new puzzle piece of T with l_i must be a digon l_il_{i+1} for $1 \leq i \leq |P| - 1$, and $l_{|P|}$ must be r . Then S is a sphere, a contradiction. Therefore, there is no $T \in \mathbb{T}_{\mathcal{S}}$ with CIV.

Case 3. $g \geq 1, |M| = 2, P = \emptyset$. Suppose that there is a $T \in \mathbb{T}_{\mathcal{S}}$ with CIV. By $P = \emptyset$, all puzzle pieces on \mathcal{S} are triangles. Moreover, \mathcal{S} has one of the following:

- (3a) one boundary component with two marked points;
- (3b) two boundary components with one marked point.

In the case (3a), a triangle of T with a boundary segment must contain a loop, a contradiction. In the case (3b), there is at least one $\gamma \in \mathbb{A}_{\mathcal{S}}$ connecting the distinct marked points. Then each triangle of T with γ must contain a boundary segment as follows:




It reduces to (3a), hence a contradiction.


Case 4. $g \geq 1, |M \setminus P| \geq 3$. We construct $T \in \mathbb{T}_{\mathcal{S}}$ with CIV for $|M| = 3$ and $P = \emptyset$, thus for all cases by Proposition 2.13. In which case, \mathcal{S} has one of the following:

- (4a) one boundary component with three marked points;
- (4b) two boundary components, one has one marked point and the other has two marked points;
- (4c) three boundary components with one marked point.


In the case (4a), we take the following triangle Δ :



We obtain a marked surface \mathcal{S}' from \mathcal{S} by cutting along the sides of Δ and removing Δ . Thus its genus is $g - 1$ and it has two boundary components with two marked points, and no punctures. We take the following triangles of \mathcal{S}' :




If $g = 1$, then they form a tagged triangulation T' of \mathcal{S}' . Considering T' as triangles on \mathcal{S} , we get $T = T' \cup \Delta \in \mathbb{T}_{\mathcal{S}}$ with CIV since their tagged arcs are not loops. If $g > 1$, then we can repeat the above process since it reduces to the case (4a) for genus $g - 1$. By induction on g , we get $T \in \mathbb{T}_{\mathcal{S}}$ with CIV. The cases (4b) and (4c) reduce to (4a) by considering the following triangles:



Case 5. $g \geq 1, |M \setminus P| = 2, |P| \geq 1$. We construct $T \in \mathbb{T}_{\mathcal{S}}$ with CIV for $|P| = 1$, thus for all cases by Proposition 2.13. In which case, \mathcal{S} has one of the following:

- (5a) one boundary component with two marked points;
- (5b) two boundary components with one marked point.


The both cases reduce to (4a) by considering the following triangles:



The cases 1 to 5 above cover all cases of \mathcal{S} with $\partial\mathcal{S} \neq \emptyset$. They imply that there is at least one $T \in \mathbb{T}_{\mathcal{S}}$ with CIV if and only if \mathcal{S} is one of the cases 1, 4, and 5. Thus the assertion holds. \square

Proof of Theorem 2.12 for wCIV. The assertions follow from (1), (2), and Proposition 2.9. \square

Example 2.14. For an annulus with exactly three marked points on the boundary, we consider the tagged triangulations



By Theorem 2.10, T does not have CIV and $\mu_3 T$ has CIV. In fact, for $\gamma \in \mu_3 T$ and any tagged arc δ ,

$$(\gamma|\delta) = A_{\gamma,\delta} = \text{Int}(\gamma, \delta).$$

On the other hand, for another loop $\rho(3)$, we have

$$(3|\rho(3)) = A_{3,\rho(3)} + B_{3,\rho(3)} = 2 + (-1) = 1, \text{ and } \text{Int}(3, \rho(3)) = A_{3,\rho(3)} = 2.$$

Thus we have $(T|\rho(3)) = (1, 1, 1) \neq (1, 1, 2) = \text{Int}(T, \rho(3))$.

3. CATEGORIFICATION OF CLUSTER ALGEBRAS

We denote by Q_0 and Q_1 the set of vertices and arrows of a quiver Q , respectively. We say that an oriented cycle of length one (resp., two) in Q is a *loop* (resp., a *2-cycle*).

3.1. Cluster algebras. We recall (skew-symmetric) cluster algebras with principal coefficients [FZ07]. For that, we need to prepare some notations. Let $n \in \mathbb{Z}_{\geq 0}$ and $\mathcal{F} := \mathbb{Q}(t_1, \dots, t_{2n})$ be the field of rational functions in $2n$ variables over \mathbb{Q} . A *seed with coefficients* is a pair (\mathbf{c}, Q) consisting of the following data:

- (1) $\mathbf{c} = (x_1, \dots, x_n, y_1, \dots, y_n)$ is a free generating set of \mathcal{F} over \mathbb{Q} .
- (2) Q is a quiver without loops nor 2-cycles such that $Q_0 = \{1, \dots, 2n\}$.

Then we refer to the tuple (x_1, \dots, x_n) as the *cluster*, to each x_i as a *cluster variable* and y_i as a *coefficient*. For a seed (\mathbf{c}, Q) with coefficients, the *mutation* $\mu_k(\mathbf{c}, Q) = (\mathbf{c}', \mu_k Q)$ at k ($1 \leq k \leq n$) is defined as follows:

- (1) $\mathbf{c}' = (x'_1, \dots, x'_n, y_1, \dots, y_n)$ is defined by

$$x_k x'_k = \prod_{(j \rightarrow k) \in Q_1} x_j \prod_{(j \leftarrow k) \in Q_1} y_{j-n} + \prod_{(j \leftarrow k) \in Q_1} x_j \prod_{(j \rightarrow k) \in Q_1} y_{j-n} \text{ and } x'_i = x_i \text{ if } i \neq k,$$

where $x_{n+1} = \dots = x_{2n} = 1 = y_{1-n} = \dots = y_0$.

- (2) $\mu_k Q$ is the quiver obtained from Q by the following steps:
 - (a) For any path $i \rightarrow k \rightarrow j$, add an arrow $i \rightarrow j$.
 - (b) Reverse all arrows incident to k .
 - (c) Remove a maximal set of disjoint 2-cycles.

We remark that μ_k is an involution, that is, we have $\mu_k \mu_k(\mathbf{c}, Q) = (\mathbf{c}, Q)$. Moreover, it is elementary that $\mu_k(\mathbf{c}, Q)$ is also a seed with coefficients.

For a quiver Q without loops nor 2-cycles such that $Q_0 = \{1, \dots, n\}$, we obtain the quiver \hat{Q} (resp., \check{Q}) from Q by adding vertices $\{1', \dots, n'\}$ and arrows $\{i \rightarrow i' \mid 1 \leq i \leq n\}$ (resp., $\{i \leftarrow i' \mid 1 \leq i \leq n\}$). We fix a seed $(\mathbf{c} = (x_1, \dots, x_n, y_1, \dots, y_n), \hat{Q})$ with coefficients, called the *initial seed*. We also call the tuple (x_1, \dots, x_n) the *initial cluster*, and each x_i the *initial cluster variable*.

Definition 3.1. The *cluster algebra* $\mathcal{A}(Q) = \mathcal{A}(\mathbf{c}, \hat{Q})$ with *principal coefficients* for the initial seed (\mathbf{c}, \hat{Q}) is a \mathbb{Z} -subalgebra of \mathcal{F} generated by all cluster variables and coefficients obtained from (\mathbf{c}, \hat{Q}) by sequences of mutations.

We denote by $\text{cl-var } Q$ (resp., *cluster* Q) the set of cluster variables (resp., clusters) of $\mathcal{A}(Q)$. One of the remarkable properties of cluster algebras with principal coefficients is the strongly Laurent phenomenon as follows.

Proposition 3.2 ([FZ07, Proposition 3.6]). *Every $x \in \text{cl-var } Q$ is expressed by a Laurent polynomial of $x_1, \dots, x_n, y_1, \dots, y_n$*

$$x = \frac{F(x_1, \dots, x_n, y_1, \dots, y_n)}{x_1^{d_1(x)} \dots x_n^{d_n(x)}},$$

where $d_i(x) \in \mathbb{Z}$ and $F(x_1, \dots, x_n, y_1, \dots, y_n) \in \mathbb{Z}[x_1, \dots, x_n, y_1, \dots, y_n]$ is not divisible by any x_i .


Definition 3.3. Using the notations in Proposition 3.2, we call $d(x) := (d_i(x))_{1 \leq i \leq n}$ the *denominator vector* of x , and $f(x) := (f_i(x))_{1 \leq i \leq n}$ the *f-vector* of x , where $f_i(x)$ is the maximal degree of y_i in the F -polynomial $F(1, \dots, 1, y_1, \dots, y_n) \in \mathbb{Z}[y_1, \dots, y_n]$ of x .

We consider the following property of $\mathcal{A}(Q)$ or Q :


$$(d=f) \quad d(x) = f(x) \text{ for any } x \in \text{cl-var } Q \setminus \{\text{initial cluster variables}\}.$$

In [FZ07, Conjecture 7.17], it was conjectured that any cluster algebra satisfies (d=f), but it does not hold in general. In fact, the following counterexample was given in [FK10, Subsection 6.4].

Example 3.4. Let Q be a quiver



and we compute the following sequence of mutations:



where we put each cluster variable or coefficient on the corresponding vertex and the non-initial cluster variables are expressed by Laurent polynomials

$$x'_3 = \frac{y_3 x_1 + x_2}{x_3}, x'_2 = \frac{y_3 x_1^2 + x_1 x_2 + y_2 y_3 x_3}{x_2 x_3}, x'_1 = \frac{y_3 x_1^2 + (y_1 y_2 y_3^2 + 1) x_1 x_2 + y_1 y_2 y_3 x_2^2 + y_2 y_3 x_3}{x_1 x_2 x_3}.$$

Then we have that $d(x'_1) = (1, 1, 1) \neq (1, 1, 2) = f(x'_1)$.

Remark 3.5. In this paper, we only consider skew-symmetric cluster algebras. However, we can also consider the property (d=f) for skew-symmetrizable cluster algebras. In fact, it is known that (d=f) holds for skew-symmetrizable cluster algebras of rank 2 [FZ07, Corollary 10.10] or finite type [Gyo21, Theorem 1.8].

Finally, we prepare the following notation.

Definition 3.6 ([Kel17]). A *reddening sequence* of Q is a sequence μ of mutations such that $\mu(\hat{Q}) = \hat{Q}$.

Note that a reddening sequence is also called a green-to-red sequence [Mul16], and Q admits a reddening sequence if and only if a seed (c, Q) is injective-reachable [Qin17].

3.2. Categorification of cluster algebras. We briefly recall a categorification of cluster algebras. Let Q be a quiver without loops nor 2-cycles and W a non-degenerate potential of Q (see [DWZ08]). In particular, such W exists if K is uncountable [DWZ08, Corollary 7.4]. The mutation of quivers with potentials is defined to be compatible with mutations of quivers, that is, $\mu_k(Q, W) = (\mu_k Q, W')$ for $k \in Q_0$ and some non-degenerate potential W' of $\mu_k Q$. Moreover, (Q, W) defines the Jacobian algebra $J = J(Q, W)$, the Ginzburg differential graded algebra $\Gamma = \Gamma_{Q, W}$, and the generalized cluster category

$\mathcal{C} = \mathcal{C}_{Q,W}$ (see [Ami09, DWZ08, Gin06, Kel08, Kel11]). The category \mathcal{C} is a triangulated category with a rigid object Γ , that is $\text{Hom}_{\mathcal{C}}(\Gamma, \Sigma\Gamma) = 0$, and $\text{End}_{\mathcal{C}}(\Gamma)^{\text{op}} \simeq J$, where Σ is the suspension functor.

For the idempotent e_i associated with $i \in Q_0$, let $\Gamma_i = \Gamma e_i$ and $\Gamma'_i = \Gamma_{\mu_k(Q,W)} e_i$. We take the cone Γ_k^* of the morphism in \mathcal{C}

$$\Gamma_k \rightarrow \bigoplus_{(k \leftarrow i) \in Q_1} \Gamma_i$$

whose components are given by the right multiplications of arrows. We say that $\mu_k \Gamma := (\Gamma/\Gamma_k) \oplus \Gamma_k^*$ is the *mutation of Γ at $k \in Q_0$* . Moreover, there is a triangle equivalence

$$\mathcal{C}_{\mu_k(Q,W)} \rightarrow \mathcal{C}$$

which sends Γ'_i to Γ_i for $i \neq k$ and Γ'_k to Γ_k^* [KY11, Theorem 3.2] (see also [Pla11b, Subsection 2.6]). Then we consider sequences of mutations from Γ as follows: For $k_1, \dots, k_j \in Q_0$, there is a triangle equivalence from $\mathcal{C}_{\mu_{k_j} \dots \mu_{k_1}(Q,W)}$ to \mathcal{C} given by the sequence of the above triangle equivalences

$$\mathcal{C}_{\mu_{k_j} \dots \mu_{k_1}(Q,W)} \rightarrow \dots \rightarrow \mathcal{C}_{\mu_{k_1}(Q,W)} \rightarrow \mathcal{C},$$

and $\mu_{k_j} \dots \mu_{k_1} \Gamma$ is defined by the image of $\Gamma_{\mu_{k_j} \dots \mu_{k_1}(Q,W)}$. Note that $\mu_{k_j} \dots \mu_{k_1} \Gamma$ is also rigid. We denote by $\text{rigid}^{\Gamma} \mathcal{C}$ the set of isomorphism classes of rigid objects in \mathcal{C} obtained from Γ by sequences of mutations, and by $\text{irigid}^{\Gamma} \mathcal{C}$ the set of isomorphism classes of indecomposable direct summands of $\Gamma' \in \text{rigid}^{\Gamma} \mathcal{C}$. Let $\text{mod} J$ be the category of finite dimensional left J -modules.

Proposition 3.7 ([Pla11b, Lemma 3.2 and Subsection 3.3]). *There is a full subcategory \mathcal{D} of \mathcal{C} including all objects in $\text{irigid}^{\Gamma} \mathcal{C}$ with an equivalence of categories*

$$\text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma, -) : \mathcal{D}/(\Gamma) \xrightarrow{\sim} \text{mod End}_{\mathcal{C}}(\Sigma^{-1}\Gamma)^{\text{op}} \simeq \text{mod} J,$$

where (Γ) is the ideal of morphisms of \mathcal{C} which factor through Γ .

Let $P_i = J e_i$ for $i \in Q_0$. By $\text{Hom}_{\mathcal{C}}(\Gamma_i, \Sigma\Gamma) = 0$ and Proposition 3.7, we have

$$\begin{aligned} \text{Hom}_J(P_i, \text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma, X)) &\simeq \text{Hom}_J(\text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma, \Sigma^{-1}\Gamma_i), \text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma, X)) \\ &\simeq \text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma_i, X). \end{aligned}$$

In particular, taking $X = \Sigma^{-1}\Gamma_i$, we have

$$(3.1) \quad \text{End}_J(P_i) \simeq \text{End}_{\mathcal{C}}(\Gamma_i).$$

The following is a main result of the categorification of cluster algebras.

Theorem 3.8. (1) [Pla11b, Theorem 4.1][IKLFP13, Corollary 3.5] *There is a bijection*

$$C_{\Gamma} : \text{rigid}^{\Gamma} \mathcal{C} \leftrightarrow \text{cl-var } Q$$

which sends Γ_i to the initial cluster variable x_i for $i \in Q_0$ and induces a bijection

$$C_{\Gamma} : \text{rigid}^{\Gamma} \mathcal{C} \leftrightarrow \text{cluster } Q$$

which sends Γ to the initial cluster and commutes with mutations.

(2) [DWZ10, Proposition 3.1][Pla11a, Subsection 3.3] *For $X \in \text{irigid}^{\Gamma}$, we have*

$$f(C_{\Gamma}(X)) = \underline{\dim} \text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma, X),$$

where the right side is the dimension vector of the J -module $\text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma, X)$.

Proposition 3.7 and Theorem 3.8 give the map in the introduction

$$M = M_{Q,W} := \text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma, C_{\Gamma}^{-1}(-)) : \text{cl-var } Q \setminus \{\text{initial cluster variables}\} \rightarrow \text{mod } J,$$

and $f(x) = \underline{\dim} M(x)$ for $x \in \text{cl-var } Q \setminus \{\text{initial cluster variables}\}$. This implies the following.

Corollary 3.9. *The quiver Q satisfies $(d=\dim)$ if and only if it satisfies $(d=f)$.*

On the other hand, if J is finite dimensional, then \mathcal{C} is a Hom-finite Krull-Schmidt 2-Calabi-Yau triangulated category and Γ is a cluster tilting object [Ami09, Theorem 2.1]. From this viewpoint, it is important to check the finite dimensionality of J and one way to do it is given as a consequence of [Pla11b, Lemma 2.17].

Proposition 3.10 ([BDP14, Proposition 8.1]). *If Q admits a reddening sequence, then $J(Q, W')$ is finite dimensional for any non-degenerate potential W' of Q .*

4. CLUSTER ALGEBRAS FROM TRIANGULATED SURFACES

4.1. Cluster algebras from triangulated surfaces. Let \mathcal{S} be a marked surface. To $T \in \mathbb{T}_{\mathcal{S}}$, we associate a quiver Q_T with $(Q_T)_0 = T$ and whose arrows correspond with angles between tagged arcs of T as in Table 3. We obtain a quiver Q_T without loops nor 2-cycles from Q_T by removing 2-cycles. This construction commutes with flips and mutations, that is, $\mu_{\gamma}Q_T = Q_{\mu_{\gamma}T}$ for $\gamma \in T$ [FST08, Proposition 4.8 and Lemma 9.7]. This implies that a quiver Q is obtained from Q_T by a sequence of mutations if and only if $Q = Q_{T'}$ for some $T' \in \mathbb{T}_{\mathcal{S}}^T$.




TABLE 3. The quiver Q_{Δ} associated with each puzzle piece Δ

Example 4.1. For marked surfaces in Theorem 1.4(2) (or Theorem 2.12(2)), it is easy to see the associated quivers as in Table 4. In particular, \mathcal{S} is a polygon without puncture (resp., with exactly one puncture) if and only if there is a $T \in \mathbb{T}_{\mathcal{S}}$ such that Q_T is a Dynkin quiver of type A (resp., D). Moreover, \mathcal{S} is an annulus with exactly two marked points (resp., a torus with exactly one puncture) if and only if all associated quivers are Kronecker (resp., Markov) quivers. These immediately give Corollary 1.5 from Theorem 1.4(2).

The associated cluster algebra $\mathcal{A}(Q_T)$ has the following properties.

Theorem 4.2. *Let $T \in \mathbb{T}_{\mathcal{S}}$.*

- (1) [FST08, Theorem 7.11][FT18, Theorem 6.1] *There is a bijection*

$$x_T : \mathbb{A}_{\mathcal{S}}^T \leftrightarrow \text{cl-var } Q_T.$$

Moreover, it induces a bijection

$$x_T : \mathbb{T}_{\mathcal{S}}^T \leftrightarrow \text{cluster } Q_T$$

which sends T to the initial cluster of $\mathcal{A}(Q_T)$ and commutes with flips and mutations.

- (2) [FZ07, (7.7)][FST08, Lemma 9.20] *For $\gamma \in \mathbb{A}_{\mathcal{S}}^T$, we have*

$$d(x_T(\gamma)) = (T|\gamma).$$

- (3) [Yur19, Theorem 7] *For $\gamma \in \mathbb{A}_{\mathcal{S}}^T$, we have*

$$f(x_T(\gamma)) = \text{Int}(T, \gamma).$$

T				
Q_T	$1 \leftarrow 2 \leftarrow 3$	$\begin{array}{c} 1 \\ \downarrow \\ 2 \\ \swarrow \quad \searrow \\ 3 \quad 4 \end{array}$	$1 \rightleftharpoons 2$	$\begin{array}{ccc} & 3 & \\ \swarrow & & \searrow \\ 1 & \rightleftharpoons & 2 \end{array}$

TABLE 4. The quiver Q_T associated with a tagged triangulation T of each marked surface in Theorem 1.4(2)

(4) [Mil17, Theorem 1.1] *If \mathcal{S} is not a closed surface with exactly one puncture, then Q_T admits a reddening sequence.*

By Theorem 4.2, we can apply the results in Section 2 to cluster algebras. In particular, we get Theorems 1.3 and 1.4 from the following corollary.


Corollary 4.3. *For $T \in \mathbb{T}_{\mathcal{S}}$, it has wCIV if and only if Q_T satisfies (d=dim).*

Proof. The assertion follows from Corollary 3.9 and Theorem 4.2. □

Proof of Theorem 1.3. The assertion follows from Theorems 2.10, 2.11, and Corollary 4.3. □

Proof of Theorem 1.4. The assertions follows from Theorem 2.12 and Corollary 4.3. □

Example 4.4. For the tagged triangulation T in Example 2.14, Q_T is the quiver in Example 3.4. Moreover, the sequence of mutations in Example 3.4 corresponds with the sequence of flips



In particular, we have $\rho(3) = 1'$ and $x_T(i') = x'_i$ for $i \in \{1, 2, 3\}$, where x'_i is the non-initial cluster variable of $\mathcal{A}(Q_T)$ appearing in Example 3.4. Then we can see $d(x_T(1')) = (T|1') = (1, 1, 1) \neq (1, 1, 2) = f(x_T(1')) = \text{Int}(T, 1')$.

4.2. Categorification on triangulated surfaces. Let \mathcal{S} be a marked surface and $T \in \mathbb{T}_{\mathcal{S}}$. The following property is given as a consequence of the classification of non-degenerate potentials of Q_T studied in [GGS15, GLFS16, GLFMO22, LF09, LF16b, Lad12] (see also [LF16a]). It also follows from Proposition 3.10 and Theorem 4.2(4).

Theorem 4.5 (Finite dimensionality). *If \mathcal{S} is not a closed surface with exactly one puncture, then $J(Q_T, W)$ is finite dimensional for any non-degenerate potential W of Q_T .*

We remark that if \mathcal{S} is a closed surface with exactly one puncture, then there is a non-degenerate potential W such that $J(Q, W)$ is infinite dimensional [GLFS16, Proof of Proposition 9.13].

Let W be a non-degenerate potential of Q_T . In the rest of this paper, we denote $J = J(Q_T, W)$, $\Gamma = \Gamma_{Q_T, W}$, $\mathcal{C} = \mathcal{C}_{Q_T, W}$, $P_\gamma = Je_\gamma$, and $\Gamma_\gamma = \Gamma e_\gamma$ for $\gamma \in T = (Q_T)_0$. By Theorems 3.8 and 4.2, we have the bijection

$$(4.1) \quad \mathsf{X}_T := \mathsf{C}_\Gamma^{-1} x_T : \mathbb{A}_{\mathcal{S}} \leftrightarrow \text{irigid}^\Gamma \mathcal{C}$$

which sends $\gamma \in T$ to Γ_γ and $\text{Int}(T, \delta) = \underline{\dim} \text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma, \mathbf{X}_T(\delta))$. The following theorem was given in [BQ15] under the assumption that J is finite dimensional, but the assumption automatically holds by Theorem 4.5.

Theorem 4.6 ([BQ15, Theorem 3.8]). *If \mathcal{S} is not a closed surface with exactly one puncture, then we have $\mathbf{X}_T(\rho(\gamma)) = \Sigma \mathbf{X}_T(\gamma)$ for $\gamma \in \mathbb{A}_{\mathcal{S}}$.*

We are ready to give equivalences including Theorem 1.6.

Theorem 4.7. *If \mathcal{S} is not a closed surface with exactly one puncture, then the following are equivalent:*

- (1) Q_T satisfies $(d=\dim)$;
- (2) $\text{End}_{\mathcal{C}}(\Gamma_\gamma) \simeq K$ for any $\gamma \in T$;
- (3) $\text{End}_J(P_\gamma) \simeq K$ for any $\gamma \in T$.

Proof. Under the assumption, CIV and wCIV are equivalent. Then (1) is equivalent to $\text{Int}(\gamma, \rho^{-1}(\gamma)) = 1$ for any $\gamma \in T$ by Theorem 2.10 and Corollary 4.3. By (4.1) and Theorem 4.6, we have

$$\begin{aligned} \text{Int}(\gamma, \rho^{-1}(\gamma)) &= \dim \text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma_\gamma, \mathbf{X}_T(\rho^{-1}(\gamma))) \\ &= \dim \text{Hom}_{\mathcal{C}}(\Sigma^{-1}\Gamma_\gamma, \Sigma^{-1} \mathbf{X}_T(\gamma)) \\ &= \dim \text{End}_{\mathcal{C}}(\Gamma_\gamma). \end{aligned}$$

Therefore, (1) and (2) are equivalent. The equivalence between (2) and (3) follows from (3.1). \square

We also consider Corollary 1.5 from the viewpoint of cluster categories. The following theorem was also given in [Yur20] under the assumption that J is finite dimensional, but the assumption automatically holds by Theorem 4.5.

Theorem 4.8 ([Yur20, Theorem 1.4]). *If \mathcal{S} is not a closed surface with exactly one puncture, then all indecomposable rigid objects in \mathcal{C} are contained in $\text{irigid}^{\Gamma} \mathcal{C}$.*

We add one condition to Corollary 1.5 under the assumption that \mathcal{S} is not closed surface with exactly one puncture.

Proposition 4.9. *If \mathcal{S} is not closed surface with exactly one puncture, then the following are equivalent:*

- (1) Q_T satisfies $(d=\dim)$ for any $T \in \mathbb{T}_{\mathcal{S}}$;
- (2) Q_T is either Dynkin type or a Kronecker quiver for any $T \in \mathbb{T}_{\mathcal{S}}$;
- (3) $\text{End}_{\mathcal{C}}(X) \simeq K$ for any indecomposable rigid object $X \in \mathcal{C}$.

Proof. By the assumption and Theorem 2.7, we have $\mathbb{T}_{\mathcal{S}} = \mathbb{T}_{\mathcal{S}}^T$. Then the equivalence between (1) and (2) is just Corollary 1.5.

For a sequence μ of mutations, the definition of mutations of Γ gives that


$$\text{End}_{\mathcal{C}}(\mu\Gamma) \simeq \text{End}_{\mathcal{C}_{\mu(Q_T, W)}}(\Gamma_{\mu(Q_T, W)}).$$

Therefore, by Theorem 4.7, $\mu(Q_T) = Q_{\mu T}$ satisfies $(d=\dim)$ if and only if $\text{End}_{\mathcal{C}_{\mu(Q_T, W)}}(X) \simeq K$ for any indecomposable direct summand X of $\Gamma_{\mu(Q_T, W)}$, which is also equivalent to $\text{End}_{\mathcal{C}}(Y) \simeq K$ for any indecomposable direct summand Y of $\mu\Gamma$. Thus the equivalence between (1) and (3) follows from Theorem 4.8. \square

5. EXAMPLE

We consider an example which is not covered by [BMR09]. For an annulus with three marked points on the boundary and one puncture, we consider the following tagged triangulation and the associated

quiver:




We take a non-degenerate potential W_T of Q_T given in [LF09] (for a puncture p , we take the parameter $x_p = -1$) as follows:

$$W_T = ba_6a_2 + ca_3a_5 - a_6a_5a_4a_3a_2a_1.$$

Then the associated Jacobian algebra $J(T) := J(Q_T, W_T)$ is the quotient of the path algebra KQ_T by the ideal generated by

$$a_6a_2, a_3a_5, a_6a_5a_4a_3a_2, a_3a_2a_1a_6a_5, \\ ba_6 - a_1a_6a_5a_4a_3, a_2b - a_5a_4a_3a_2a_1, ca_3 - a_4a_3a_2a_1a_6, a_5c - a_2a_1a_6a_5a_4.$$

The indecomposable projective module $J(T)e_3$ is described by




where all paths from the top vertex to each vertex coincide in $J(T)$ and the paths for all vertices form a basis of $J(T)e_3$. In particular, the dimension vector is given by the numbers of tagged arcs appearing in the diagram, that is,

$$\underline{\dim} J(T)e_3 = (2, 2, 4, 2, 2).$$

Therefore, since $\dim \text{End}_{J(T)}(J(T)e_3) = 4 \neq 1$, Q_T does not satisfy (d=dim) by Theorem 4.7. Note that the dimension vector is also given by $\text{Int}(T, \rho^{-1}(3))$ in the same way as the proof of Theorem 4.7.

Next, we consider the tagged triangulation $\mu_3 T$ and the associated quiver:




We take a non-degenerate potential $W_{\mu_3 T}$ of $Q_{\mu_3 T}$ as follows:

$$W_{\mu_3 T} = \gamma_2 \beta_1 \alpha_4 + \beta_2 \gamma_1 \alpha_2 + \alpha_4 \alpha_3 \alpha_2 \alpha_1.$$

Note that the quivers with potential $(Q_{\mu_3 T}, W_{\mu_3 T})$ and $\mu_3(Q_T, W_T)$ are right-equivalent [LF09, Theorem 30]. Then the associated Jacobian algebra $J(\mu_3 T) := J(Q_{\mu_3 T}, W_{\mu_3 T})$ is the quotient of the path algebra $KQ_{\mu_3 T}$ by the ideal generated by

$$\begin{aligned} &\alpha_4 \gamma_2, \beta_1 \alpha_4, \alpha_2 \beta_2, \gamma_1 \alpha_2, \alpha_4 \alpha_3 \alpha_2, \alpha_2 \alpha_1 \alpha_4 \\ &\gamma_2 \beta_1 - \alpha_3 \alpha_2 \alpha_1, \beta_2 \gamma_1 - \alpha_1 \alpha_4 \alpha_3. \end{aligned}$$

All indecomposable projective $J(\mu_3 T)$ -modules are described as follows:



We can see that $\dim \text{End}(J(\mu_3 T)e_i) = 1$ for any $1 \leq i \leq 5$ and $Q_{\mu_3 T}$ satisfies (d=dim) by Theorem 4.7.

Acknowledgements. The author would like to thank Pierre-Guy Plamondon for helpful comments on Proposition 3.10. He was JSPS Overseas Research Fellow and supported by JSPS KAKENHI Grant Numbers JP20J00410, JP21K13761.

REFERENCES

[Ami09] Claire Amiot. Cluster categories for algebras of global dimension 2 and quivers with potential. In *Annales de l'institut Fourier*, volume 59, pages 2525–2590, 2009.

[BDP14] Thomas Brüstle, Grégoire Dupont, and Matthieu Pérotin. On maximal green sequences. *International Mathematics Research Notices*, 2014(16):4547–4586, 2014.

[BMR09] Aslak Bakke Buan, Robert J Marsh, and Idun Reiten. Denominators of cluster variables. *Journal of the London Mathematical Society*, 79(3):589–611, 2009.

- [BQ15] Thomas Brüstle and Yu Qiu. Tagged mapping class groups: Auslander–reiten translation. *Mathematische Zeitschrift*, 279(3-4):1103–1120, 2015.
- [CC06] Philippe Caldero and Frédéric Chapoton. Cluster algebras as hall algebras of quiver representations. *Commentarii Mathematici Helvetici*, 81(3):595–616, 2006.
- [CK06] Philippe Caldero and Bernhard Keller. From triangulated categories to cluster algebras ii. In *Annales scientifiques de l’Ecole normale supérieure*, volume 39, pages 983–1009, 2006.
- [DWZ08] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky. Quivers with potentials and their representations i: Mutations. *Selecta Mathematica*, 14(1):59–119, 2008.
- [DWZ10] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky. Quivers with potentials and their representations ii: applications to cluster algebras. *Journal of the American Mathematical Society*, 23(3):749–790, 2010.
- [FG06] Vladimir Fock and Alexander Goncharov. Moduli spaces of local systems and higher teichmüller theory. *Publications Mathématiques de l’IHÉS*, 103:1–211, 2006.
- [FK10] Changjian Fu and Bernhard Keller. On cluster algebras with coefficients and 2-calabi-yau categories. *Transactions of the American Mathematical Society*, 362(2):859–895, 2010.
- [FST08] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces. part i: Cluster complexes. *Acta Mathematica*, 201(1):83–146, 2008.
- [FT18] Sergey Fomin and Dylan Thurston. *Cluster algebras and triangulated surfaces Part II: Lambda lengths*, volume 255. American Mathematical Society, 2018.
- [FZ02] Sergey Fomin and Andrei Zelevinsky. Cluster algebras i: foundations. *Journal of the American Mathematical Society*, 15(2):497–529, 2002.
- [FZ07] Sergey Fomin and Andrei Zelevinsky. Cluster algebras iv: coefficients. *Compositio Mathematica*, 143(1):112–164, 2007.
- [GGS15] Christof Geiss and Raúl González-Silva. Tubular jacobian algebras. *Algebras and Representation Theory*, 18(1):161–181, 2015.
- [Gin06] Victor Ginzburg. Calabi-yau algebras. *arXiv preprint math/0612139*, 2006.
- [GLFMO22] Jan Geuenich, Daniel Labardini-Fragoso, and José Luis Miranda-Olvera. Quivers with potentials associated to triangulations of closed surfaces with at most two punctures. *Séminaire Lotharingien de Combinatoire*, B84c:21 pp, 2022.
- [GLFS16] Christof Geiß, Daniel Labardini-Fragoso, and Jan Schröer. The representation type of jacobian algebras. *Advances in Mathematics*, 290:364–452, 2016.
- [Gyo21] Yasuaki Gyoda. Relation between f-vectors and d-vectors in cluster algebras of finite type or rank 2. *Annals of Combinatorics*, 25(3):573–594, 2021.
- [Hub06] Andrew Hubery. Acyclic cluster algebras via ringel-hall algebras. *Preprint, January*, 2006.
- [IKLFP13] Giovanni Cerulli Irelli, Bernhard Keller, Daniel Labardini-Fragoso, and Pierre-Guy Plamondon. Linear independence of cluster monomials for skew-symmetric cluster algebras. *Compositio Mathematica*, 149(10):1753–1764, 2013.
- [Kel08] Bernhard Keller. Calabi-yau triangulated categories. *Trends in representation theory of algebras and related topics*, pages 467–489, 2008.
- [Kel11] Bernhard Keller. Deformed calabi–yau completions. *Journal für die reine und angewandte Mathematik*, 654:125–180, 2011. with an appendix by Michel Van den Bergh.
- [Kel17] Bernhard Keller. Quiver mutation and combinatorial dt-invariants. *arXiv preprint arXiv:1709.03143*, 2017.
- [KY11] Bernhard Keller and Dong Yang. Derived equivalences from mutations of quivers with potential. *Advances in Mathematics*, 226(3):2118–2168, 2011.
- [Lad12] Sefi Ladkani. On jacobian algebras from closed surfaces. *arXiv preprint arXiv:1207.3778*, 2012.
- [LF09] Daniel Labardini-Fragoso. Quivers with potentials associated to triangulated surfaces. *Proceedings of the London Mathematical Society*, 98(3):797–839, 2009.
- [LF16a] Daniel Labardini-Fragoso. On triangulations, quivers with potentials and mutations. *Mexican Mathematicians Abroad*, 657:103, 2016.
- [LF16b] Daniel Labardini-Fragoso. Quivers with potentials associated to triangulated surfaces, part iv: removing boundary assumptions. *Selecta Mathematica*, 22(1):145–189, 2016.
- [Mar16] Bruno Martelli. An introduction to geometric topology. *arXiv preprint arXiv:1610.02592*, 2016.
- [Mil17] Matthew R Mills. Maximal green sequences for quivers of finite mutation type. *Advances in Mathematics*, 319:182–210, 2017.
- [Mul16] Greg Muller. The existence of a maximal green sequence is not invariant under quiver mutation. *The Electronic Journal of Combinatorics*, 23(2):P2–47, 2016.
- [Pla11a] Pierre-Guy Plamondon. Cluster algebras via cluster categories with infinite-dimensional morphism spaces. *Compositio Mathematica*, 147(6):1921–1954, 2011.
- [Pla11b] Pierre-Guy Plamondon. Cluster characters for cluster categories with infinite-dimensional morphism spaces. *Advances in Mathematics*, 227(1):1–39, 2011.
- [Qin17] Fan Qin. Triangular bases in quantum cluster algebras and monoidal categorification conjectures. *Duke Math. J.*, 166(12):2337–2442, 2017.

- [QZ17] Yu Qiu and Yu Zhou. Cluster categories for marked surfaces: punctured case. *Compositio Mathematica*, 153(9):1779–1819, 2017.
- [RS17] Nathan Reading and Salvatore Stella. Initial-seed recursions and dualities for d-vectors. *Pacific Journal of Mathematics*, 293(1):179–206, 2017.
- [Yur19] Toshiya Yurikusa. Combinatorial cluster expansion formulas from triangulated surfaces. *The Electronic Journal of Combinatorics*, 26(2):P2–33, 2019.
- [Yur20] Toshiya Yurikusa. Density of g-vector cones from triangulated surfaces. *International Mathematics Research Notices*, 2020(21):8081–8119, 2020.

LABORATOIRE DE MATHÉMATIQUES DE VERSAILLES, UVSQ, CNRS, UNIVERSITÉ PARIS-SACLAY, 78035 VERSAILLES, FRANCE AND MATHEMATICAL INSTITUTE, TOHOKU UNIVERSITY, SENDAI 980-8578, JAPAN
Email address: toshiya.yurikusa.d8@tohoku.ac.jp